

Assembly, configuration, and break-up history of Rodinia: A synthesis

Z.X. Li^{a,g,*}, S.V. Bogdanova^b, A.S. Collins^c, A. Davidson^d, B. De Waele^a, R.E. Ernst^{e,f},
I.C.W. Fitzsimons^g, R.A. Fuck^h, D.P. Gladkochubⁱ, J. Jacobs^j, K.E. Karlstrom^k,
S. Lu^l, L.M. Natapov^m, V. Peaseⁿ, S.A. Pisarevsky^a, K. Thrane^o, V. Vernikovsky^p

^a Tectonics Special Research Centre, School of Earth and Geographical Sciences, The University of Western Australia, Crawley, WA 6009, Australia

^b Department of Geology, Lund University, Solvegatan 12, 223 62 Lund, Sweden

^c Continental Evolution Research Group, School of Earth and Environmental Sciences, University of Adelaide, Adelaide, SA 5005, Australia

^d Geological Survey of Canada (retired), 601 Booth Street, Ottawa, Canada K1A 0E8

^e Ernst Geosciences, 43 Margrave Avenue, Ottawa, Canada K1T 3Y2

^f Department of Earth Sciences, Carleton U., Ottawa, Canada K1S 5B6

^g Tectonics Special Research Centre, Department of Applied Geology, Curtin University of Technology, GPO Box U1987, Perth, WA 6845, Australia

^h Universidade de Brasília, 70910-000 Brasília, Brazil

ⁱ Institute of the Earth's Crust SB RAS, Lermontova Street, 128, 664033 Irkutsk, Russia

^j Department of Earth Science, University of Bergen, Allegaten 41, N-5007 Bergen, Norway

^k Department of Earth and Planetary Sciences, Northrop Hall University of New Mexico, Albuquerque, NM 87131, USA

^l Tianjin Institute of Geology and Mineral Resources, CGS, No. 4, 8th Road, Tianjin 300170, China

^m GEMOC ARC Key Centre, Department of Earth and Planetary Sciences, Macquarie University, Sydney, NSW 2109, Australia

ⁿ Department of Geology and Geochemistry, Stockholm University, 10691 Stockholm, Sweden

^o Geological Institute, University of Copenhagen, Øster Voldgade 10, 1350 Copenhagen, Denmark

^p Institute of Geology & Mineralogy of Russian Academy of Sciences, Koptyug Avenue 3, Novosibirsk 630090, Russia

Received 27 June 2006; received in revised form 1 December 2006; accepted 23 April 2007

Abstract

This paper presents a brief synthesis of the current state of knowledge on the formation and break-up of the early-Neoproterozoic supercontinent Rodinia, and the subsequent assembly of Gondwanaland. Our discussions are based on both palaeomagnetic constraints and on geological correlations of basement provinces, orogenic histories, sedimentary provenance, the development of continental rifts and passive margins, and the record of mantle plume events.

Rodinia assembled through worldwide orogenic events between 1300 Ma and 900 Ma, with all, or virtually all, continental blocks known to exist at that time likely being involved. In our preferred Rodinia model, the assembly process features the accretion or collision of continental blocks around the margin of Laurentia. Like the supercontinent Pangaea, Rodinia lasted about 150 million years after complete assembly. Mantle avalanches, caused by the sinking of stagnated slabs accumulated at the mantle transition zone surrounding the supercontinent, plus thermal insulation by the supercontinent, led to the formation of a mantle superswell (or superplume) beneath Rodinia 40–60 million years after the completion of its assembly. As a result, widespread continental rifting occurred between ca. 825 Ma and 740 Ma, with episodic plume events at ca. 825 Ma, ca. 780 Ma and ca. 750 Ma.

Like its assembly, the break-up of Rodinia occurred diachronously. The first major break-up event occurred along the western margin of Laurentia (present coordinates), possibly as early as 750 Ma. Rifting between the Amazonia craton and the southeastern margin of Laurentia started at approximately the same time, but only led to break-up after ca. 600 Ma. By this time most of the western Gondwanan continents had joined together, although the formation of Gondwanaland was not complete until ca. 530 Ma.

© 2007 Elsevier B.V. All rights reserved.

Keywords: Rodinia; Supercontinent; Neoproterozoic; Mesoproterozoic; Grenvillian; Continental rift; Mantle plume; Gondwanaland

* Corresponding author. Present address: Department of Applied Geology, Curtin University of Technology, GPO Box U1987, Perth, WA 6845, Australia.
Tel.: +61 8 9266 2453; fax: +61 8 9266 3153.

E-mail address: z.li@curtin.edu.au (Z.X. Li).

1. Introduction

Valentine and Moores (1970) were probably the first to recognise that a supercontinent, comprising just about all continents on Earth, existed towards the end of the Precambrian. They suggested that the break-up of the supercontinent, which they called Pangaea I, by the Cambrian led to a divergence of environments and characteristics amongst the daughter continents, and a sudden prominence of shallow, nutrient-rich shelves and coastal areas, all factors conducive to the diversification of life forms on Earth. This late-Precambrian supercontinent was later renamed Rodinia (McMenamin and McMenamin, 1990) from the Russian word ‘rodit’ meaning ‘to beget’ or ‘to give birth’. McMenamin and McMenamin (1990) considered Rodinia to have been the supercontinent that spawned all subsequent continents, while “the edges (continental shelves) of Rodinia were the cradle of the earliest animals” (McMenamin and McMenamin, 1990, p. 95).

The concept of the supercontinent Rodinia attracted much attention in 1991, when three researchers (Moores, 1991; Dalziel, 1991; Hoffman, 1991) published geological evidence for the assembly and break-up of Rodinia, with some of its daughter continents forming Gondwanaland. Common to their propositions is the connection between western Laurentia, Australia and East Antarctica (commonly known as the southwest U.S.–East Antarctic, or SWEAT, connection; Moores, 1991), which follows an earlier suggestion by Bell and Jefferson (1987) based on stratigraphic correlations and palaeomagnetic constraints across the Pacific, and by Eisbacher (1985) based on stratigraphic correlation alone.

An explosion of new data and ideas occurred in the following years, and, in order to better coordinate the global efforts in testing the Rodinia hypothesis, a UNESCO-IGCP project (No. 440, 1999–2004) was established to investigate the formation, configuration, and break-up of Rodinia, and to construct an interpretative Geodynamic Map of the Rodinia supercontinent (the Rodinia Map hereafter; Appendix I, see hard copy in the print version of this volume; also available online). Although few still doubt the existence of a late Precambrian supercontinent, there is still no consensus regarding the number of participating cratons, their relative configuration within the supercontinent and the chronology and mode of assembly and break-up of the supercontinent. In this paper we provide an overview of evidence for and against major Rodinian reconstructions, including those in the Rodinia Map (Appendix I). We discuss some current ideas regarding the formation of Rodinia and processes that led to its break-up, and present an animation for the evolution of Rodinia from 1100 Ma till the formation of Gondwanaland at 530 Ma (Appendix II, available online).

The challenges to reconstructing the history of Rodinia include inadequate high-quality geological, geochronological and palaeomagnetic data, multiple possible interpretations for each data set, and uncertainties in fundamental assumptions such as the application of modern-style plate tectonics to late-Precambrian time and that the geomagnetic field was a geocentric axial dipole field (an assumption that underlines interpretations of palaeomagnetic data). In this paper we assume that modern-style plate tectonics, with the possible

complication of true polar wander (TPW), apply to late Precambrian (e.g., Stern, 2005), and that the geomagnetic field was dominantly a geocentric axial dipole field at that time (Evans, 2006). We emphasise the importance of considering multiple lines of evidence in testing any reconstruction, because only through such an approach can the potentially large number of solutions be reduced to the most likely scenarios (Fig. 1a). We strive to be as objective as possible when discussing alternative interpretations, at the same time trying to make a self-consistent synthesis. We emphasise that not every opinion expressed in this paper is agreed on by all co-authors.

2. Major inter-continental connections proposed for Rodinia using multidisciplinary evidence

Because Laurentia is flanked by Neoproterozoic passive margins, it is commonly regarded as being at the centre of Rodinia assembly and break-up (e.g., Hoffman, 1991). We will evaluate the various continental connections around the margin of Laurentia proposed for Rodinia time.

2.1. Continents facing the present western and northern margins of Laurentia: Australia–East Antarctica, South China, or Siberia?

2.1.1. The SWEAT hypothesis

The question of which continent(s) used to lie adjacent to the present western margin of Laurentia during the existence of Rodinia has been central to the Rodinia debate, and the topic remains controversial today. Among the competing models, the earliest and perhaps the best-known model is the SWEAT hypothesis (Fig. 2a). The model builds on the similar Neoproterozoic stratigraphy of the western margin of Laurentia and eastern Australia as recognised earlier by Eisbacher (1985) and Bell and Jefferson (1987), and elaborated later by Young (1992) and Rainbird et al. (1996) among others. Three landmark papers in 1991 (Moores, 1991; Dalziel, 1991; Hoffman, 1991) also used basement outcrops as piercing points that may correlate from one continent to the other, such as the correlation of Grenville-age (ca. 1300–1000 Ma) orogenic belts, to argue that the SWEAT connection probably existed as early as 1900 Ma ago and lasted until mid to late Neoproterozoic times. Moores (1991) implied that the united northern and western Australian craton might not have become part of the SWEAT connection until its collision with the Gawler craton along the late Mesoproterozoic Albany–Fraser–Musgrave belt. Powell et al. (1993) confirmed that a SWEAT connection was palaeomagnetically permissible for the time interval of ca. 1050–720 Ma based on the contemporary palaeomagnetic database, but would have to have broken apart by 580 Ma (or >650 Ma as pointed out earlier by Van der Voo et al., 1984, on palaeomagnetic grounds).

Closer scrutiny of the crustal provinces and basin provenance analysis, however, have shown that geological discontinuities exist across the SWEAT connection. In particular, isotopic and geochemical mapping along the Transantarctic Mountains has revealed that Mesoproterozoic basement provinces as young as ca. 1000 Ma along the continental margin truncate the Lau-




Fig. 1. (a) A schematic diagram illustrating the importance of taking a multidisciplinary and global approach in reconstructing palaeogeography in relation to the evolution of a Precambrian supercontinent (when biogeography is of limited value). The line from each research aspect symbolises an infinite number of possible solutions, with concentration of their intersections giving the most likely solution. (b) Present locations and major crustal elements of the Precambrian continental blocks.

rentian crustal provinces (Borg and DePaolo, 1994). To make the SWEAT connection viable, these authors suggested that those basement blocks were probably exotic in origin and were accreted to the East Antarctic margin after the opening of the proto-Pacific Ocean. Although this may be so for some of the terranes along the Cambrian Ross–Delamerian Orogen (Stump, 1995), there is no geological evidence for the docking of terranes like the Beardmore terrane, which would have to have occurred in a very narrow time window between the break-up

of the SWEAT connection (ca. 750 Ma, see later discussions) and the development of the Beardmore Group passive-margin deposits no later than 668 ± 1 Ma (Goodge et al., 2002).

There are other lines of evidence that do not support the SWEAT connection. (1) There is a lack of continuation of the ca. 1400 Ma (1500–1350 Ma) transcontinental magmatic province of southern Laurentia (e.g., Nyman et al., 1994; Van Schmus et al., 1996) into the Transantarctic Mountains, although one could argue that they could be beneath the ice farther inland (Goodge

et al., 2002). (2) There are mismatches in the Neoproterozoic mantle plume record across the SWEAT connection (Park et al., 1995; Wingate et al., 1998; Li et al., 1999; see more discussion in Section 2.1.2). (3) The Belt Basin of western North America and the possibly overlying Buffalo Hump Formation

(the Deer Trail Group) require a western clastic source with rocks of 1786–1642 Ma, 1600–1590 Ma and 1244–1070 Ma, much of which cannot easily be identified in East Antarctica in the SWEAT configuration (Ross et al., 1992; Ross and Villeneuve, 2003). (4) The originally proposed continuation of




Fig. 2. Palaeomagnetic examinations of alternative configurations of continents west and north of Laurentia. Palaeopoles used are listed in Table 1, and Euler rotation parameters given in Appendix III (available online). (a) In the SWEAT fit (after Moores, 1991; Dalziel, 1991; Hoffman, 1991) palaeopoles from Laurentia (grey diamonds) and Australia (purple diamonds) do not merge until ca. 1050 Ma. (b) In the “Missing-Link” configuration (after Li et al., 1995) ca. 820–800 Ma poles from India, Australia and South China fall close to each other but the ca. 750 Ma poles are scattered. No dated ca. 820–800 Ma pole is available from Laurentia for a comparison. (c) In the AUSWUS configuration (Karlstrom et al., 1999; Burrett and Berry, 2000) ca. 1200 Ma poles from Laurentia and Australia fall >30° apart, whereas some of their 1100–1050 Ma poles follow the same apparent polar wander path (APWP). (d) In the AUSMEX configuration (Wingate et al., 2002) five of the seven ca. 1200–1050 Ma poles from Australia fall away from coeval Laurentian poles. (e) Siberia placed against Laurentia following Sears and Price (1978, 2000), in which the Siberian poles fall away from the Laurentia poles. (f) Siberia–Laurentia configuration following Pisarevsky et al. (2008), in which ca. 1050–970 Ma Siberian poles follow the same APWP as the Laurentian poles. In this figure and Figs. 5–7, all continents were rotated to Laurentia which is at its 900 Ma palaeolatitude. Latitude and longitude lines are shown in 30° intervals. Geotectonic polygon data are shown following the legend for the Rodinia Map (as in Figs. 1b and 8 and Appendix I).




Fig. 2. (Continued).

the Grenville belt into the Coats Land and Dronning Maud Land of East Antarctica (Moores, 1991; Dalziel, 1991) was later proved untenable both palaeomagnetically (e.g., Gose et al., 1997) and geologically (e.g., Jacobs et al., 2003a,b). Coats Land was more likely part of Laurentia at 1110 Ma (e.g., Gose et al., 1997), but became connected with Kalahari when Kalahari collided with Laurentia at 1090–1060 Ma, and remained connected until the break-up of Gondwanaland (e.g., Jacobs et al., 2003a,b, 2008).

There is also a growing body of evidence suggesting the possible presence of Grenvillian orogenic belts along both the western margin of Laurentia and eastern Australia. On the Australia–East Antarctica side, apart from the possible presence of late Mesoproterozoic basement provinces along the Transantarctic Mountains (Borg and DePaolo, 1994), Berry et al. (2005) recently reported a 1287 ± 18 Ma metamorphic age (monazite U–Th–Pb) from King Island northwest of Tasmania (#1 in Fig. 2a), and Fioretti et al. (2005) published a 1119 ± 9 Ma age for quartz syenite at the South Tasman Rise. Like the basement for the Transantarctic Mountains, an allochthonous origin for northwestern Tasmania and King Island is possible, but the following observations argue against such a possibility: (a) the palaeomagnetic record shows that northwestern Tasmania was already part of cratonic Australia by the Late Cambrian (Li et al., 1997), and there is no prior compressional event recognised in the region that reflects terrane amalgamation between the time of Rodinia break-up (ca. 750 Ma, see Section 3.2) and the Late Cambrian; (b) the Neoproterozoic tectonostratigraphy of northwestern Tasmania and King Island can be well correlated with that of the Adelaide Fold Belt (Calver and Walter, 2000; Holm et al., 2003; Li, 2001; Z.X. Li et al., 2003b). A possible Grenvillian province has also been identified in northern Queensland (#2 in Fig. 2a; Blewett et al., 1998; Hutton et al., 1996), largely based on detrital zircon ages.

A number of studies reporting Grenvillian orogenic events along the western margin of Laurentia have yet to attract much attention. Perhaps most significant are the 1090–1030 Ma metamorphic ages (titanite U–Pb) from 1468 ± 2 Ma mafic sills in the Belt–Purcell Supergroup (#3 in Fig. 2a; Anderson and Davis, 1995). In the Mackenzie Mountains (#4 in Fig. 2a), there is evidence for an east–west compressional event (the Corn Creek Orogeny) that occurred after 1033 Ma but before 750 Ma (Thorkelson, 2000; Thorkelson et al., 2005). There is also a less-well defined 1175–1100 Ma recrystallisation age from a granite clast brought up from the basement in a diatreme (Jefferson and Parrish, 1989).

If there were indeed Grenvillian orogenic belts between Laurentia and Australia–East Antarctica, it would suggest that even if a SWEAT connection existed, it would have to have been Grenvillian or later.

Palaeomagnetic data permit a SWEAT connection after ca. 1050 Ma (Powell et al., 1993), but such a connection is not possible at ca. 1200 Ma (Fig. 2a; Pisarevsky et al., 2003a) although the reliability of the ca. 1200 Ma pole for Australia needs further verification. This scenario is consistent with pre-Grenvillian geological mismatches discussed above. However, we do not preclude the possible existence of a SWEAT-like link between Australia and Laurentia during late-Paleoproterozoic (ca. 1800–1600 Ma), as suggested by Idnurm and Giddings (1995) and Betts and Giles (2006).

2.1.2. The “Missing-Link” model

The “Missing-Link” model proposed by Li et al. (1995) has the South China Block sitting between Australia–East Antarctica and Laurentia in Rodinia, serving as the “missing-link” between the two continents (Fig. 2b). The model was initially developed in view of: (a) mismatches in the crustal provinces of Australia–East Antarctica and Laurentia (see discussions in

Table 1

Selected palaeomagnetic poles used for reconstructing the late-Mesoproterozoic to Neoproterozoic global palaeogeography

| Rock unit | Age (Ma) | Pole | | A ₉₅ (°) | Reference |
|-------------------------------------|---------------------|------|-------|---------------------|---|
| | | (°N) | (°E) | | |
| Laurentia | | | | | |
| Long Range dykes ^a | 620–610 | 19 | 355 | 18 | Murthy et al. (1992); Kamo and Gower (1994) |
| Franklin dykes | 723 + 4/–2 | 5 | 163 | 5 | Heaman et al. (1992); Park (1994) |
| Natkusiak Formation | 723 + 4/–2 | 6 | 159 | 6 | Palmer et al. (1983); Heaman et al. (1992) |
| Kwagunt Formation | 742 ± 6 | 18 | 166 | 7 | Weil et al. (2004) |
| Tsezotene sills and dykes | 779 ± 2 | 2 | 138 | 5 | Park et al. (1989); LeCheminant and Heaman (1994) |
| Wyoming dykes | 782 ± 8; 785 ± 8 | 13 | 131 | 4 | Harlan et al. (1997) |
| Galeros Formation | 780–820 | –2 | 163 | 6 | Weil et al. (2004) |
| Haliburton intrusions A | 980 ± 10; 1000–1030 | –36 | 143 | 10 | Buchan and Dunlop (1976); Warnock et al. (2000) |
| Chequamegon sandstone | ~1020 ^b | –12 | 178 | 5 | McCabe and Van der Voo (1983) |
| Jacobsville sandstone J (A+B) | ~1020 ^b | –9 | 183 | 4 | Roy and Robertson (1978) |
| Freda sandstone | 1050 ± 30 | 2 | 179 | 4 | Henry et al. (1977); Wingate et al. (2002) |
| Nonesuch shale | 1050 ± 30 | 8 | 178 | 4 | Henry et al. (1977); Wingate et al. (2002) |
| Lake Shore Traps | 1087 ± 2 | 22 | 181 | 5 | Diehl and Haig (1994); Davis and Paces (1990) |
| Portage Lake volcanics | 1095 ± 2 | 27 | 181 | 2 | Halls and Pesonen (1982); Davis and Paces (1990) |
| Upper North Shore volcanics | 1097 ± 2 | 32 | 184 | 5 | Halls and Pesonen (1982); Davis and Green (1997) |
| Logan sills R | 1109 + 4/–2 | 49 | 220 | 4 | Halls and Pesonen (1982); Davis and Sutcliffe (1985) |
| Abitibi dykes | 1141 ± 2 | 43 | 209 | 14 | Ernst and Buchan (1993) |
| Wind River, Gr. B | 1300–1500 | 22 | 209 | 9 | Harlan et al. (2003a,b) |
| Mackenzie dolerite dykes | 1267 ± 2 | 4.0 | 190.0 | 5.0 | Buchan and Halls (1990) |
| Nain anorthosite | 1320–1290 | 12.0 | 210.0 | 3.0 | From Buchan et al. (2000) |
| Mistastin complex | ~1420 | –1.0 | 201.0 | 8.0 | From Buchan et al. (2000) |
| Laramie complex and Sherman granite | ~1434 | –7.0 | 215.0 | 4.0 | From Buchan et al. (2000) |
| Harp Lake complex | ~1450 | 2.0 | 206.0 | 4.0 | From Buchan et al. (2000) |
| Michikamau anorthosite pluton | ~1460 | –2.0 | 218.0 | 5.0 | From Buchan et al. (2000) |
| Molson dykes, component A | ~1820–1720 | 15.4 | 263.5 | 4.0 | From Buchan et al. (2000) |
| Baltica | | | | | |
| Hunnedalen dykes | >848 | –41 | 222 | 10 | Walderhaug et al. (1999) |
| Egersund-Ogna anorthosite | ~900 | –42 | 200 | 9 | Brown and McEnroe (2004) |
| Egersund anorthosite | 929–932 | –44 | 214 | 4 | Stearn and Piper (1984); Torsvik and Eide (1998) |
| Hakefjorden | 916 ± 11 | 5 | 249 | 4 | Stearn and Piper (1984); Scherstén et al. (2000) |
| Göteborg-Slussen | 935 ± 3 | –7 | 242 | 12 | Pisarevsky and Bylund (2006) |
| Dalarna dykes | 946 ± 1 | 5 | 239 | 15 | Bylund and Elming (1992), Söderlund et al. (2005) |
| Karlshamn-Fäjö dykes | 946–954 | 2 | 242 | 30 | Patchett and Bylund (1977), Söderlund et al. (2004) |
| Nilstorp dyke | 966 ± 2 | 9 | 239 | 8 | Patchett and Bylund (1977), Söderlund et al. (2004) |
| Pyätteryd amphibolite | 933–945 | –43 | 214 | 11 | Pisarevsky and Bylund (1998); Wang et al. (1996); Wang and Lindh (1996) |
| Känna gneiss | 948–974 | –50 | 225 | 17 | Pisarevsky and Bylund (1998); Wang et al. (1996); Wang and Lindh (1996) |
| Gällared amphibolite | 956? | –46 | 214 | 19 | Pisarevsky and Bylund (1998); Möller and Söderlund (1997) |
| Gällared granite-gneiss | 980–990 | –44 | 224 | 6 | Pisarevsky and Bylund (1998); Möller and Söderlund (1997) |
| Bamble intrusions | 1100–1040 | 3 | 217 | 15 | Torsvik and Eide (1998); Brown and McEnroe (2004) |
| Laanila dolerite | 1045 ± 50 | –2 | 212 | 15 | Mertanen et al. (1996) |
| Mean Jotnian dolerite intrusions | ~1265 | 4.0 | 158.0 | 4.0 | From Buchan et al. (2000) |
| Åland quartz porphyry dykes | 1571 ± 20; 1571 ± 9 | 12.0 | 182.0 | 7.0 | From Buchan et al. (2000) |
| Åland dolerite dykes | 1577 ± 12 | 28.0 | 188.0 | 9.0 | From Buchan et al. (2000) |
| Subjotnian quartz porphyry dykes | ~1630 | 29.0 | 177.0 | 6.0 | From Buchan et al. (2000) |
| Shoksha Formation, Vepsian Group | ~1780 | 39.7 | 221.1 | 4.0 | From Buchan et al. (2000) |

Table 1 (Continued)

| Rock unit | Age (Ma) | Pole | | A ₉₅ (°) | Reference |
|---|--------------------|-------|-------|---------------------|--|
| | | (°N) | (°E) | | |
| India | | | | | |
| Mahe dykes, Seychelles ^c | 748–755 | 80 | 79 | 11 | Torsvik et al. (2001b) |
| Malani Igneous Suite | 751–771 | 68 | 88 | 8 | Torsvik et al. (2001a) |
| Harohalli dykes | 814 ± 34 | 27 | 79 | 18 | Radhakrishna and Mathew (1996) |
| Wajrakarur kimberlites | 1079? | 45 | 59 | 11 | Miller and Hargraves (1994) |
| Australia | | | | | |
| Elatina Formation | 600–620 | 51 | 157 | 2 | Embleton and Williams (1986) |
| Elatina Formation | 600–620 | 54 | 147 | 1 | Schmidt et al. (1991) |
| Elatina Formation | 600–620 | 52 | 167 | 11 | Schmidt and Williams (1995) |
| Elatina Formation | 600–620 | 39 | 186 | 9 | Sohl et al. (1999) |
| Yaltipena Formation | 620–630 | 44 | 173 | 8 | Sohl et al. (1999) |
| Mundine Well dykes | 755 ± 3 | 45 | 135 | 4 | Wingate and Giddings (2000) |
| Walsh Tillite | 750–770 | 22 | 102 | 14 | Li (2000) |
| Hussar Formation | 800–760 | 62 | 86 | 10 | Pisarevsky et al. (2007) |
| Browne Formation | 830–800 | 45 | 142 | 7 | Pisarevsky et al. (2007) |
| Wooltana Formation | ~820 | 62 | 322 | 17 | McWilliams and McElhinny (1980) |
| Kulgera dykes | 1054 ± 13 | −8 | 75 | 6 | Schmidt et al. (2005) |
| Bangemall Basin sills | 1070 ± 6 | 34 | 95 | 8 | Wingate et al. (2002) |
| Stuart dykes | 1076 ± 33 | 10 | 262 | 10 | Idnurm and Giddings (1988) |
| Lakeview dolerite | 1116 ± 12 | 10 | 311 | 17 | Tanaka and Idnurm (1994) |
| Bremer Bay & Whalebone Pt | ~1200 | 74 | 304 | 13 | Pisarevsky et al. (2003a) |
| Mt. Barren Group | 1205 ± 40 | 47 | 347 | 13 | Pisarevsky et al. (2003a), Dawson et al. (2003) |
| Fraser dyke | 1212 ± 10 | 56 | 3260 | 5 | Pisarevsky et al. (2003a) |
| Congo | | | | | |
| Mbozi Complex, Tanzania | 755 ± 25 | 46 | 325 | 9 | Meert et al. (1995); Evans (2000) |
| Gagwe lavas, Tanzania | 795 ± 7 | 25 | 93 | 10 | Meert et al. (1995); Deblond et al. (2001) |
| São Francisco | | | | | |
| Ilheus dykes | 1011 ± 24 | 30 | 100 | 4 | D’Agrella-Filho et al. (1990); Renne et al. (1990) |
| Oliveira dykes, normal | ~1035 ^b | 16 | 107 | 8 | D’Agrella-Filho et al. (1990); Renne et al. (1990) |
| Itaju de Colonia | ~1055 ^b | 8 | 111 | 10 | D’Agrella-Filho et al. (1990); Renne et al. (1990) |
| Oliveira dykes (rever.) | 1078 ± 18 | −10 | 100 | 9 | D’Agrella-Filho et al. (1990); Renne et al. (1990) |
| Kalahari ^d | | | | | |
| Ritscherflya Supergroup (rotated to Kalahari) | 1130 ± 12 | 61 | 29 | 4 | Powell et al. (2001) |
| Umkondo Igneous Province | 1105 ± 5 | 66 | 37 | 3 | Powell et al. (2001); Wingate (2001) |
| Kalkpunt Formation, | ~1065? | 57 | 3 | 7 | Briden et al. (1979); Powell et al. (2001) |
| Central Namaqua | ~1030–1000 | 8 | 330 | 10 | Onstott et al. (1986); Robb et al. (1999) |
| Siberia | | | | | |
| Ust-Kirba sediments and sills | ~980 | −8 | 183 | 10 | Pavlov et al. (2002) |
| Kandyk Formation | ~1000 | −3 | 177 | 4 | Pavlov et al. (2002); Rainbird et al. (1998) |
| Ignikan Formation | ~1010 | −16 | 201 | 4 | Pavlov et al. (2000) |
| Nelkan Formation | ~1020 | −14 | 219 | 6 | Pavlov et al. (2000) |
| Milkon Formation | ~1025 | −6 | 196 | 4 | Pavlov et al. (2000) |
| Kumahinsk Formation | ~1030 | −14 | 201 | 7 | Pavlov et al. (2000) |
| Malgina Formation | 1043 ± 14 | −22 | 226 | 7 | Osipova in Smethurst et al. (1998); Ovchinnikova et al. (2001) |
| Malgina Formation | 1043 ± 14 | −25 | 231 | 3 | Gallet et al. (2000); Ovchinnikova et al. (2001) |
| Chieress dyke | 1384 ± 2 | 4 | 258 | 6.6 | Ernst et al. (2000) |
| Kuonamka dykes | 1503 ± 5 | 6 | 234 | 22 | Ernst et al. (2000) |
| North China | | | | | |
| Dongjia Formation, Lushan | ~650 | −60.8 | 97.4 | 6.7 | From Zhang et al. (2006) |
| Mean poles | ~700 | −42.9 | 107.0 | 5.7 | From Zhang et al. (2006) |
| Nanfen Formation | 800–780 | −16.5 | 121.1 | 11.1 | From Zhang et al. (2006) |
| Cuizhuang/Sanjiaotang Formations | 950 | −41.0 | 44.8 | 11.3 | From Zhang et al. (2006) |
| Tieling Formation | 1100 | 2.2 | 163.6 | 25.3 | From Zhang et al. (2006) |
| Baicaoping Formation | 1200 | −43.0 | 143.8 | 11.1 | Zhang et al. (2006) |

Table 1 (Continued)

| Rock unit | Age (Ma) | Pole | | A_{95} (°) | Reference |
|--------------------------------|--------------|-------|------|--------------|--------------------------|
| | | (°N) | (°E) | | |
| Yunmengshan Formation | 1260 | −60.6 | 87.0 | 3.7 | Zhang et al. (2006) |
| Yangzhuang/Wumishan Formations | 1300 | −17.2 | 45.0 | 5.5 | From Zhang et al. (2006) |
| Yangzhuang Formation | ~1350 | −17.3 | 3.5 | 5.7 | Wu et al. (2005) |
| Taihang dyke swarm | 1769.1 ± 2.5 | −36.0 | 67.0 | 2.8 | Halls et al. (2000) |
| Tarim | | | | | |
| Aksu dykes | 807 ± 12 | 19 | 128 | 6 | Chen et al. (2004) |
| Baiyixi Formation | ~740 | 17 | 194 | 4 | Huang et al. (2005) |
| South China | | | | | |
| Doushantuo carbonates | 584 ± 26 | 6 | 197 | 6 | Macouin et al. (2004) |
| Liantuo Formation | 748 ± 12 | 4 | 161 | 13 | Evans et al. (2000) |
| Xiaofeng dykes | 802 ± 10 | 14 | 91 | 11 | Li et al. (2004) |
| Amazonia | | | | | |
| Nova Floresta | ~1200 | 25 | 165 | 6 | Tohver et al. (2002) |
| Oaxaquia | | | | | |
| Oaxaca anorthosite | ~950 | 47 | 267 | 23 | Ballard et al. (1989) |

Abbreviations: SA, South Australia; NT, Northern Territory; WA, Western Australia.

^a Recalculated by Hodych et al. (2004).

^b Age based on APWP interpolation.

^c Rotated to India 28° counterclockwise around the pole of 25.8°N, 330°E (Torsvik et al., 2001b).

^d For ages see Powell et al. (2001) and references therein.

the previous section); (b) similarities in the Neoproterozoic stratigraphy of South China, southeastern Australia and western Laurentia as recognised earlier by Eisbacher (1985); (c) similarities between crustal provinces in the Cathaysia Block of southeastern South China and southern Laurentia; (d) the need for a western source region to provide the late Mesoproterozoic detrital grains in the Belt Basin. The modified reconstruction (Fig. 2b) and the schematic history for the formation and break-up of such a configuration were adapted for constructing the Geodynamic Map of Rodinia (Appendix I) and the related palaeogeographic time slices and animation. Although not a unique solution, it negates the need for matching basement geology between Australia–East Antarctica and western Laurentia prior to 1000–900 Ma (see discussions in Section 2.1.1), and is plausible in view of the currently known Neoproterozoic rift record, early Neoproterozoic mantle plume record, and palaeomagnetic constraints. Key evidence for the model is discussed below.

2.1.2.1. Cathaysia as an extension of Laurentia from ca. 1800 Ma until Rodinia time, and no connection between Australia–East Antarctica, Yangtze craton and Laurentia–Cathaysia until 1100–900 Ma. The Cathaysia Block, although poorly exposed, has a crustal composition similar to what may be expected for the western source region of the Belt Basin in southwestern Laurentia (Ross et al., 1992). It has a ca. 1830–1430 Ma basement that recorded 1300–1000 Ma metamorphism (Z.X. Li et al., 2002). Hainan Island in this block is particularly similar to the Mojave Province in that it has ca. 1430 Ma granitic intrusions, as well as synchronous intracratonic sedimentary and volcanic successions (Z.X. Li et al., 2002, and unpublished data), comparable to the 1500–1350 Ma transcontinental granite–rhyolite province of

southern Laurentia (e.g., Nyman et al., 1994). The sedimentary provenance of quartzite in the overlying post-1200 Ma succession (Li et al., unpublished data), assumed to have sourced from older Cathaysian rocks, contains populations comparable to the 1610–1490 Ma, westerly sourced non-Laurentian detrital grains reported in the Belt Basin (Ross and Villeneuve, 2003). It is thus plausible to have Cathaysia as a part of Laurentia for at least the latter part of the 1830–1000 Ma interval (Figs. 2b and 3).

According to the “Missing-Link” model, the collision between Laurentia–Cathaysia and Yangtze did not start until ca. 1140 Ma or younger at one end of the Sibao Orogen (Greentree et al., 2006), and lasted until ca. 900 Ma at the other end in South China (Z.X. Li et al., 2003a; Ling et al., 2003). This late formation of the Laurentia–South China–Australia–East Antarctica connection provides an explanation for the occurrence of Grenvillian orogenic events along western Laurentia, northern Queensland, King Island–Transantarctic Mountains, and the Albany–Fraser–Musgrave belt.

2.1.2.2. Neoproterozoic plume record. Large features related to mantle plume events, such as radiating dyke swarms or large igneous provinces, can be used for reconstructing past continental connections (e.g., Ernst et al., 1995, 2008). Park et al. (1995) were the first to apply this technique to test the SWEAT connection, but precise dating of the relevant dyke swarms revealed that their ages are too different to represent the same event: ca. 825 Ma for the Gairdner–Amata dyke swarm in central and southeastern Australia (Sun and Sheraton, 1996; Wingate et al., 1998), and ca. 780 Ma for the radiating Gunbarrel dyke swarms in western Laurentia (e.g., Harlan et al., 2003a,b). No counterpart of the Australian 825 Ma event has been reported in western Laurentia as one would expect in the SWEAT configuration. The radiating 780 Ma Gunbarrel dyke swarm in western Laurentia




Fig. 3. Tectonostratigraphic correlations between Australia, Yangtze craton, Cathaysia and Laurentia. Major glaciations: M = Marinoan; S = Sturtian; N = Nantuo; C = Changan; I = Ice Brook; R = Rapitan.

points to a plume-centre to its west (present coordinates), but in eastern Australia there is no evidence for a plume centre apart from limited volcanism in the Adelaide Rift Complex (Preiss, 2000) and minor mafic igneous rocks in northwestern Tasmania (Holm et al., 2003).

Li et al. (1999) and Z.X. Li et al. (2003b) presented evidence for South China being above a plume centre at both ca. 825 Ma and 780 Ma, which is consistent with its position in the "Missing-Link" configuration (Fig. 4). Key evidence for a ca. 825 Ma plume-head beneath South China includes the widespread occurrence of magmatism with rock types ranging from granite to mafic-ultramafic dykes and sills (implying mafic underplating and a large heat source), large-scale syn-magmatic doming, and the development of ca. 820 Ma continental rift systems (Li et al., 1999; X.H. Li et al., 2003a,b; Ling et al., 2003; Wang and Li, 2003). A similar large-scale magmatic breakout occurred at ca. 780 Ma, with some mafic dykes showing geochemical characteristics of continental flood basalts (Z.X. Li et al., 2003b; Lin et al., 2007). It should be noted that alternative geochemical interpretations exist for the 825–740 Ma magmatism in South China (e.g., arc magmatism, Zhou et al., 2002; magmatism caused by post-orogenic slab break-off, Wang et al., 2006).

2.1.2.3. Neoproterozoic rift records and glacial events. Rifting records in South China show remarkable similarities to that of eastern Australia, featuring four major episodes of magmatism and rifting in the ca. 830–700 Ma interval: ca. 820 Ma, ca. 800 Ma, ca. 780 Ma, and ca. 750–720 Ma (Powell et al., 1994;

Li et al., 1995, 1999; X.H. Li et al., 2002; Z.X. Li et al., 2003b; Preiss, 2000; Wang and Li, 2003) (Fig. 3). A rift–drift transition has been identified in the Adelaidean stratigraphy to be between the ca. 720 Ma Sturtian glacial deposits and the overlying Tapley Hill Formation (Powell et al., 1994).

There is no record of rifting along western Laurentia until after the 780 Ma Gunbarrel dyke swarm (e.g., Harlan et al., 2003a,b). Rift magmatism mainly occurred between 750 Ma and 720 Ma (e.g., Heaman et al., 1992; Ross and Villeneuve, 1997; Karlstrom et al., 2000), although evidence exists for events of younger ages (Lund et al., 2003; Fanning and Link, 2004) as in South China (e.g., Zhou et al., 2004; Zhang et al., 2005). Along with the glacial record, the post-780 Ma tectonostratigraphy of western Laurentia, eastern Australia (e.g., Young, 1992; Rainbird et al., 1996) and South China (e.g., Wang and Li, 2003) are widely regarded as correlative (Fig. 3), although such a correlation is not unique across the globe.

2.1.2.4. Palaeomagnetism. Another advantage of the "Missing-Link" model is that it does not require a common apparent polar wander path (APWP) between the relevant continents until amalgamation sometime between 1000 Ma and 900 Ma. On the other hand, the currently available palaeomagnetic data for ca. 820–800 Ma permit such an Australia–South China–Laurentia configuration at that time, although it must have broken up by ca. 750 Ma (Li et al., 2004; note the overlap between the 820 Ma and 800 Ma palaeopoles for India, Australia and South China, and the scatter in the ca. 750 Ma poles in Fig. 2b).




Fig. 4. Evidence for plume activities in the Missing-Link model: GD, Gairdner-Amata dyke swarm (Sun and Sheraton, 1996; Wingate et al., 1998); BH, Bow Hill lamprophyre dykes (Pidgeon et al., 1989); GB, Guibei mafic-ultramafic intrusions (Li et al., 1999); GUN, Gunbarrel dyke swarms (Harlan et al., 2003a,b); KD, Kangding dykes and mafic intrusions (Z.X. Li et al., 2003b; Lin et al., 2007); T, Mafic and granitic intrusions in Tasmania (Holm et al., 2003).

However, the configuration in Fig. 2b is not unique either geologically or palaeomagnetically. An alternative position for South China northeast of Australia satisfies the ca. 750 Ma poles (Evans et al., 2000), but contradicts the ca. 800 Ma poles (Li et al., 2004). Positions to the northwest or west of Australia, and adjacent to the northern margin of Greater India, have also been proposed (Evans et al., 2000; Jiang et al., 2003; Li et al., 2004; Yang et al., 2004), but the geological matches in these configurations are not as strong as in the “Missing-Link” configuration discussed above. Also, by removing South China from the “Missing-Link” configuration, one would have to find other explanations for the geological and palaeomagnetic mismatches between Australia–East Antarctica and Laurentia as discussed in Section 2.1.1.

2.1.3. The AUSWUS and AUSMEX connections

AUSWUS (Australia–Southwest US)-like fit (Fig. 2c) was first suggested by Brookfield (1993) based on matching linear fractures supposedly formed during the break-up of Rodinia along the margins of eastern Australian craton and western Laurentia. However, the lineaments along the eastern margin of the Australian craton were later shown to be no older than ca. 600 Ma (Direen and Crawford, 2003). The AUSWUS connection was later revived by Karlstrom et al. (1999) and Burrett and Berry (2000), mostly through matching basement provinces and sedimentary provenances between Australia and southwestern Laurentia (also see Blewett et al., 1998; Berry et al., 2001; Ross and Villeneuve, 2003; Wade et al., 2006). Despite its merits as stated by its proponents, there are noticeable difficulties in this model (Fig. 2c), namely:

- (1) The truncation of the Albany-Fraser-Musgrave belt against cratonic western Laurentia, and difficulties in explaining the scattered evidence of Grenvillian metamorphism in western Laurentia, northern Queensland, Tasmania and the Ross Orogen (see Section 2.1.1).
- (2) The lack of a prominent ca. 1400 Ma granite–rhyolite province in Australia as is present in southern Laurentia.
- (3) The lack of a ca. 825 Ma plume record in Laurentia as a counterpart to the Gairdner-Amata dyke swarm in Australia (Wingate et al., 1998), and a lack of ca. 780 Ma plume record in northern Australia, where one would expect to find a plume-head for the 780 Ma Gunbarrel radiating dyke swarm in western Laurentia in such a configuration.
- (4) The significantly younger starting age of continental rifting in western Laurentia (<750 Ma) compared with eastern Australia (ca. 825 Ma).
- (5) The ca. 1200 Ma palaeomagnetic misfit (Pisarevsky et al., 2003a), although the ca. 1100–1050 Ma palaeopoles mostly agree rather well (Fig. 2c).

Based on a new 1070 Ma palaeomagnetic pole from the Bange-mall Basin sills in the Edmund Fold Belt of Western Australia, Wingate et al. (2002) argued that previous poles of the same age from central Australia (poles marked as 1076 Ma and 1054 Ma in Fig. 2b and c) are unreliable. Using only their own pole, they argued that if Australia was connected to Laurentia at ca. 1070 Ma, the connection could only have been between northern Queensland of Australia and Mexico of southern Laurentia (the AUSMEX fit; Fig. 2d). However, subsequent work on the 1070 Ma rocks in central Australia reaffirmed the reliability of the central Australian poles (Schmidt et al., 2005). Therefore it appears probable that either the 1070 Ma Bange-mall Basin sills, or the central Australian data, have suffered vertical-axis rotations, during either the ≥ 900 Ma (^{40}Ar – ^{39}Ar cooling ages) Edmundian Orogeny in the southern Capricorn Orogen (Occhipinti and Reddy, 2005) in the case of the Bange-mall sills, or the late Neoproterozoic Petermann or Phanerozoic Alice Springs orogenies in central Australia. The geological merit of the AUSMEX fit is yet to be demonstrated. Nonetheless, a new palaeomagnetic pole from the 800 Ma to 760 Ma oriented

cores from the Lancer-1 drill hole in the Officer Basin of south Central Australia is more compatible with the ~ 780 Ma Laurentian poles in the AUSMEX configuration than in any of the abovementioned configurations (Pisarevsky et al., 2007).

As mentioned earlier, ca. 1200 Ma poles do not permit either an AUSWUS or an AUSMEX fit at that time (Pisarevsky et al., 2003a; Fig. 2c and d).

2.1.4. Siberia–Laurentia connection

Although there is a consensus that Siberia was likely connected to Laurentia for much of the Proterozoic, opinions differ regarding how they were connected. The dominant view is that Siberia was adjacent to northern Laurentia, but different models show its position in a range of orientations (Hoffman, 1991; Condie and Rosen, 1994; Pelechaty, 1996; Frost et al., 1998; Rainbird et al., 1998; Ernst et al., 2000; Vernikovskiy and Vernikovskaya, 2001; Metelkin et al., 2005a; see review by Pisarevsky et al., 2003b). A vastly different position for Siberia was proposed by Sears and Price (1978, 2000) with Siberia placed west of the present western margin of Laurentia. However, this latter fit is not supported by palaeomagnetic data (Fig. 2e) which show that Siberian poles fall far away from coeval palaeopoles for Laurentia in such a configuration. The configuration we adopted for the Rodinia Map follows that of Pisarevsky et al. (2008). It is largely based on existing palaeomagnetic constraints for the ~ 1050 – 970 Ma interval between Siberia and Laurentia (Fig. 2f, which shows that Siberian poles generally follow the APWP of Laurentia for the 1043– 980 Ma interval). Detailed discussions for the various models are given in Pisarevsky et al. (2008).

Palaeomagnetic data place Siberia at some distance from the northern Laurentian margin, allowing space for other Precambrian blocks (e.g., Arctica of Zonenshain et al., 1990; Vernikovskiy, 1997; Vernikovskiy et al., 2003). The relative distance between Siberia and Laurentia explains the lack of any counterpart for the Mackenzie large igneous event in Siberia. Sklyarov et al. (2003), Metelkin et al. (2005c) and Gladkochub et al. (2006) reported mafic intrusions in southern Siberia with ^{40}Ar – ^{39}Ar ages around ~ 740 Ma that may be correlated with the Franklin igneous event in northern Laurentia associated with the Neoproterozoic opening of the Palaeo-Asian Ocean. This event broadly coincides with the accumulation of the Karagas Group in south Siberia, which has been interpreted as a passive margin succession (Pisarevsky and Natapov, 2003; Vernikovskiy et al., 2003; Gladkochub et al., 2006). This earlier break-up model contrasts with the Early Cambrian break-up suggested by Pelechaty (1996) based on tectonostratigraphic analyses, with the latter model being challenged by both Rainbird and de Freitas (1997) and Khudoley (1997).

2.2. Continents along the present eastern margin of Laurentia: Baltica, Amazonia and West Africa cratons

The relative position of Baltica to Laurentia in Rodinia is probably one of the least controversial. Although geological correlations suggest that the two blocks could have been together as long ago as ca. 1800 Ma (e.g., Gorbatschev and Bogdanova,




Fig. 5. Positions of Baltica, Amazonia and West Africa in Rodinia, with palaeomagnetic South poles showing data from Baltica (green poles) merge with that of Laurentia (black poles) by ca. 1000 Ma. Orange dots represent ca. 1500–1350 Ma intracratonic magmatism.

1993; Karlstrom et al., 2001; see also Zhao et al., 2002 for a review) or even before (e.g., Heaman, 1997; Bleeker and Ernst, 2006), palaeomagnetic data suggest a rather complex history between the two cratons during the Mesoproterozoic (e.g., Elming et al., 1993, 2001; Buchan et al., 2000, 2001; Pesonen et al., 2003). The configuration we adapted in the Rodinia Map (Appendix I; Fig. 5) follows that of Hoffman (1991; see also Bogdanova et al., 2008; Pease et al., 2008), which is a geologically based reconstruction but is also supported by a common APWP between the two cratons for the ~ 1000 – 900 Ma interval (Elming et al., 1993; Hyodo et al., 1993; Weil et al., 1998; Pisarevsky et al., 2003b; Fig. 5). It is noted that, although pre-1000 Ma palaeomagnetic data do not permit exactly the same configuration, they still allow the two adjacent cratons to develop correlative Proterozoic belts including the Sveconorwegian and Grenville orogens (Winchester, 1988; Gower et al., 1990; Gorbatschev and Bogdanova, 1993), and similar ca. 1500–1350 Ma intracratonic magmatism (Åhäll and Connely, 1998; Karlstrom et al., 2001; Bogdanova et al., 2001, 2008; Čečys et al., 2002; Söderlund et al., 2002, 2005; Čečys and Benn, 2007).

The position of the Amazonia craton roughly follows that of Hoffman (1991, a geologically based reconstruction) and Weil et al. (1998), (a palaeomagnetic reconstruction). Amazonia was one of the cratons that was separated from Laurentia by a Mesoproterozoic ocean, the closure of which made it part of Rodinia during the Grenvillian Orogeny (e.g., Davidson, 1995,

2008; Rivers, 1997; Loewy et al., 2003). According to this interpretation, the Rondonia-Sunsas belt in southwestern Amazonia resulted from continental collision with Laurentia between 1080 Ma and 970 Ma (e.g., Sadowski and Bettencourt, 1996; Tassinari et al., 2000 and references therein). A recent ~1200 Ma Amazonia palaeomagnetic pole reported by Tohver et al. (2002) led these authors to suggest a possible juxtaposition of western Amazonia with the Llano segment of Laurentia's Grenville orogen. This is broadly in accord with the proposed history of the Llano orogen (Mosher, 1998). Tohver et al. (2004, 2005) more recently suggested that the initial docking of Amazonia with southern Laurentia was followed by strike-slip transport of Amazonia to the northeast (present coordinates). However, such palaeomagnetic reconstructions, utilizing a single pole from each continent rather than matching relatively long segments of APWPs, have inherent longitudinal uncertainties.

Continental rifting along the eastern and southern margin of Laurentia may have started as early as ca. 750 Ma (Su et al., 1994; Aleinikoff et al., 1995; Fetter and Goldberg, 1995), although a later pulse of magmatism at ~615–570 Ma is commonly interpreted as representing the break-up and opening of the Iapetus Ocean, first between Laurentia and Baltica, and then between Laurentia and Amazonia (Cawood et al., 2001; Cawood and Pisarevsky, 2006 and references therein).

There are few constraints on the position of the West Africa craton in Rodinia. No orogen of Grenvillian age has been found in or around the craton; therefore, if it was part of Rodinia, it would likely have been part of a larger craton (e.g., together with Amazonia, similar to their connection in Gondwanaland). No palaeomagnetic data are available to verify this position. Trompette (1994, 1997) proposed the existence of a single West Africa–Amazonia–Rio de La Plata mega-craton in the Meso- and Neoproterozoic, although their relative positions cannot be defined precisely. Onstott and Hargraves (1981), on the basis of a comparison of Paleo- to Mesoproterozoic palaeomagnetic data from these two blocks, also suggested that the two cratons were together during the Proterozoic but that large shear movements occurred between them. We have used an Amazonia–West Africa fit similar to their Gondwana fit.

2.3. Congo–São Francisco, Rio de la Plata and Kalahari cratons

The positions of Congo–São Francisco and Kalahari cratons relative to Laurentia in Rodinia (Fig. 6) broadly follow that proposed by Hoffman (1991) on geological grounds, but with some additional justifications according to more recent palaeomagnetic results. The Rio de la Plata terrane is placed between the




Fig. 6. Positions of Congo, Rio de la Plata, Kalahari, India and Tarim in Rodinia. Palaeomagnetic data show that Congo-Sao Francisco did not join Laurentia until after ca. 1010 Ma. The ca. 795 Ma pole from Congo-Sao Francisco overlaps with the ca. 814 Ma pole from India in this configuration, but their ca. 755 Ma poles are widely separated (as the other ca. 755 Ma poles in Fig. 2b), suggesting this configuration was already disintegrating by that time.

Congo-São Francisco craton and Laurentia in Rodinia (e.g., Weil et al., 1998; D'Agrella-Filho et al., 2004).

The preservation of late Mesoproterozoic reworked and juvenile crust and orogenic events along the Irumide belt of the Congo-São Francisco craton (e.g., De Waele et al., 2003) and along the margin of the Rio de la Plata terrane (Fuck et al., 2008) permits them to be the continents that joined Laurentia during the Grenvillian Orogeny, although there are opinions that the Congo-São Francisco craton may not have been part of Rodinia at all (e.g., Kröner and Cordani, 2003; Pisarevsky et al., 2003b). Palaeomagnetic data indicate the presence of a large latitudinal gap (an ocean?) between the Congo-São Francisco craton and Laurentia at ca. 1100–1050 Ma (D'Agrella-Filho et al., 2004); the ≥ 1000 Ma orogenic events in the Irumide belt may indicate closure of this ocean by ca. 1000 Ma (e.g., De Waele et al., 2003, 2006a,b, 2008; De Waele, 2005; Johnson et al., 2006, 2007a).

The Kalahari craton, together with the Grunehogna terrane (e.g., Groenewald et al., 1995; Jones et al., 2003) and perhaps part of the Maud Province of East Antarctica (e.g., Fitzsimons, 2000), are placed against the Grenville belt in southern Laurentia following Hoffman (1991) and Hanson et al. (1998). Palaeomagnetic data from Kalahari and Laurentia at ca. 1110 Ma suggest that there was a latitudinal difference of ca. $30 \pm 14^\circ$ between their facing margins at that time (Powell et al., 2001; Hanson et al., 2004), but by ca. 1000 Ma the APWPs of the two continents merged together in this configuration (Fig. 6). This is consistent with metamorphic ages of as young as ca. 1000 Ma along both the Namaqua-Natal belt in southern Kalahari (e.g., Eglington and Armstrong, 2003) and the Grenville belt in Laurentia (e.g., Davidson, 1995; Rivers, 1997; Keppie et al., 2003; Weber and Hecht, 2003). However, other possible positions of Kalahari in Rodinia have also been proposed on geological and palaeomagnetic grounds (e.g., Dalziel, 1997; Pisarevsky et al., 2003b).

Like the Amazonia craton, the Congo-São Francisco, Rio de la Plata and the Kalahari cratons are believed to have broken away from Laurentia between 750 Ma (see evidence below) and the final assembling of Gondwanaland by ca. 550–520 Ma (e.g., Meert, 2003; Collins and Pisarevsky, 2005). Rift magmatism at ca. 750 Ma is present along the Grenvillian margin of Laurentia (e.g., Su et al., 1994; Fetter and Goldberg, 1995; Aleinikoff et al., 1995), the western margin of Kalahari (e.g., Frimmel et al., 1996, 2001; Hoffman et al., 1996), and the Damaran margin of the Congo-São Francisco craton (e.g., Hoffmann et al., 2004; Halverson et al., 2005). The stratigraphy of the Kalahari and Congo-São Francisco continental margins indicates rift–drift transitions immediately after ca. 750 Ma (Frimmel et al., 1996; Hoffman et al., 1996; Hoffmann et al., 2004; Halverson et al., 2005; Johnson et al., 2007b). As shown in Fig. 6, ca. 750 Ma palaeomagnetic poles from the Congo-São Francisco craton fall $>30^\circ$ away from coeval poles of Laurentia.

The position of the Congo-São Francisco and Kalahari cratons in Rodinia as shown in Fig. 6 is also consistent with the common occurrence of ca. 830–745 Ma plume-induced intraplate magmatism in these two cratons as well as coeval magmatism in adjacent India, Australia, South China and Tarim

cratons (see Section 3.2). However, no pre-750 Ma Neoproterozoic magmatism has yet been identified in southern Laurentia.

2.4. India, Madagascar, Seychelles, and the Arabian–Nubian Shield

Geological data demonstrate that northeastern Madagascar (the Antongil block) was part of the Indian craton before and during Rodinia time (e.g., Collins and Windley, 2002; Collins et al., 2003; Collins, 2006)—here we collectively call them the Indian craton. The Seychelles appears to have formed along the west margin of India during the Neoproterozoic (Torsvik et al., 2001a,b).

The position of the Indian craton in Rodinia is controversial. Earlier workers attached it to Australia and East Antarctica as in the Gondwanan configuration (e.g., Dalziel, 1991; Hoffman, 1991; Moores, 1991; Li et al., 1996; Torsvik et al., 1996; Weil et al., 1998). However, subsequent geological (Fitzsimons, 2000) and palaeomagnetic (see below) investigations have challenged such a configuration. Based on their palaeomagnetic results from the 771 Ma to 751 Ma Malani igneous suite of northwestern India, Torsvik et al. (2001a,b) suggested that India was either not part of Rodinia, or was adjacent to the northwestern margin of Australia at Rodinia time and underwent a left-lateral movement relative to Australia–East Antarctica before 535 Ma to reach its Gondwanan position. Powell and Pisarevsky (2002) supported the idea that India was probably never part of Rodinia because a ca. 810 Ma pole (Radhakrishna and Mathew, 1996) places India at a polar position when Rodinia was supposed to occupy a largely low palaeolatitude position.

In the Rodinia Map (Appendix I), the position of India between 900 Ma and 800 Ma is made following both geological evidence (e.g., plume record as described by Z.X. Li et al., 2003b) and palaeomagnetic interpretations as described by Li et al. (2004). Both India and Seychelles have Neoproterozoic bimodal magmatism similar to that in Australia and South China in age distributions. Many interpreted this to represent continental arc magmatism, partly because of the perceived “continental margin” positions of these continental blocks (e.g., Torsvik et al., 2001a,b; Tucker et al., 2001; Ashwal et al., 2002). However, their bimodal nature, the intraplate characteristics of the Malani igneous suite (e.g., Roy, 2001; Singh and Vallinayagam, 2004) and other Neoproterozoic intrusions elsewhere in India (e.g., Santosh et al., 1989), and their similar age distributions to plume-related rocks in Australia and South China, make them more likely the products of melting above Neoproterozoic plume-heads during Rodinia time (e.g., Frimmel et al., 2001; Li et al., 2001; Z.X. Li et al., 2003b).

Palaeomagnetically, a poorly dated ca. 1079 Ma pole from India (Miller and Hargraves, 1994) does not agree with coeval poles from Australia in the preferred Rodinia configuration, whereas the ca. 810 Ma pole from India (Radhakrishna and Mathew, 1996) agrees with similar-aged poles from South China (Li et al., 2004), South Australia (McWilliams and McElhinny, 1980; however, the reliability of this pole has not been demonstrated with confidence) (Fig. 2b), and the Congo-São Francisco craton (Meert et al., 1995; Deblond et al., 2001) (Fig. 6).

We thus envisage that India became part of Rodinia by ca. 900 Ma through continental collision along the ca. 990–900 Ma high-grade metamorphic Eastern Ghats belt of India and the corresponding Rayner Province in East Antarctica (e.g., Mezger and Cosca, 1999; Boger et al., 2000, 2001; Fitzsimons, 2000; Kelly et al., 2002; Fig. 6).

The ca. 770–750 Ma palaeopoles from the India craton are far removed from the ca. 755 Ma pole of the Mundine Well dykes in Australia (Wingate and Giddings, 2000) in this configuration (Fig. 2b), indicating that if the configuration is correct, India would have rifted away from Australia–East Antarctica by ca. 755 Ma.

There are few constraints on the positions of the numerous terranes constituting the East African Orogen (Stern, 1994), which includes the Arabian–Nubian Shield and the ~3000 km long Azania continent that stretches from Arabia to central Madagascar (Collins and Windley, 2002; Collins and Pisarevsky, 2005; Collins, 2006). Cox et al. (2004) and Fitzsimons and Hulscher (2005) suggest that central Madagascar was part of the Congo–São Francisco craton at ca. 1800 Ma but became an independent terrane during Rodinia time and joined India by ca. 700 Ma. Most of the terranes in the Arabian–Nubian Shield are believed to represent juvenile Neoproterozoic arcs and micro-continental fragments that accreted to western Gondwana during the Neoproterozoic (e.g., Stern, 1994; Whitehouse et al., 2001; Meert, 2003; Collins and Pisarevsky, 2005). However, the nature of widespread 850–750 Ma igneous rocks in these terranes remains controversial. Some believe they represent arc volcanism (e.g., Stern, 1994; Handke et al., 1999; Tucker et al., 2001). Others suggest that those bimodal rocks were likely formed (1) during continental extension (e.g., Kröner et al., 2000; Loizenbauer et al., 2001) with possible underplating of plume magmatism, (2) with contributions of oceanic plateaux (Stein and Goldstein, 1996), or (3) as arcs with a plume input (Teklay et al., 2002). The Neoproterozoic Arabian–Nubian Shield is thought to represent one of the most rapid episodes of predominantly juvenile crustal generation in Earth's history (Stein and Goldstein, 1996; Stein, 2003). As the age spectrum of these rocks is almost identical to those in South China, Australia and India, there could indeed have been a significant plume input as part of the Neoproterozoic Rodinia superplume activity (Li et al., 2003; Hargrove et al., 2004; see further discussions in Section 3.2).

In previous Rodinia reconstructions the Arabian–Nubian Shield was either placed adjacent to northeast Madagascar and India (e.g., Li and Powell, 2001), or adjacent, and outboard of the Congo–São Francisco and Saharan cratons (Collins and Pisarevsky, 2005). In the Rodinia Map (Appendix I and Figs. 6 and 8), these terranes are placed between India and Sahara, satisfying both the common plume records in Australia, India, the Arabian Shield, and the African cratons, and the records of arc accretion during the late Neoproterozoic assembly of Gondwanaland.

2.5. Tarim

We have positioned the Tarim craton of northwest China adjacent to northwestern Australia (Fig. 6) following the suggestion

of Li et al. (1996) that was based on tectonostratigraphic correlations, with some palaeomagnetic justifications. Minor terranes (such as the Cimmerian terranes, Metcalfe, 1996) may have existed between Tarim and Australia (Li and Powell, 2001). The geological arguments for such a configuration are: (1) the presence of late Mesoproterozoic–earliest Neoproterozoic active margins along both the present northwestern and southern margin of this Archaean–Paleoproterozoic craton (Zhang et al., 2003; Lu et al., 2008a) suggests that it did not join Australia until the beginning of the Neoproterozoic; (2) 820–750 Ma bimodal (plume-induced?) intrusions in Tarim (Chen et al., 2004; Guo et al., 2005; C.L. Zhang et al., 2006; Lu et al., 2008b) can be correlated with the ca. 820–800 Ma lamprophyre dykes and kimberlite pipes in the Kimberley craton of western Australia (Pidgeon et al., 1989), and the 755 Ma Mundine Well dyke swarm in the northwestern Pilbara craton (Wingate and Giddings, 2000) and A-type granitic magmatism in the Leeuwin Block of southwestern Australia (Collins, 2003). A noticeable difference is the lack of ≤ 750 Ma rift magmatism in western Australia, which is present in Tarim (Xu et al., 2005; Z.X. Li et al., unpublished data). This could be due to the rift margin being farther off shore in Australia, or to it being totally reworked by the Pinjarra Orogeny (Fitzsimons, 2003); (3) both the Kimberley region of northwestern Australia and Tarim had up to three Neoproterozoic glacial intervals, although precise ages of the glacial events and their lateral correlations are still debated (Li et al., 1996; Grey and Corkeron, 1998; Xiao et al., 2004); (4) Cambrian mafic volcanic units in northeastern Tarim may correlate with the Antrim Plateau Volcanics in the Kimberley (Li et al., 1996), which have been interpreted as part of the 513 Ma Kalkarindji Large Igneous Province in central and northern Australia (Hanley and Wingate, 2000; Glass and Phillips, 2006).

Two Neoproterozoic palaeomagnetic poles have recently been reported from the Tarim craton: one from ca. 800 Ma mafic dykes (Chen et al., 2004), and the other from a ca. 750–730 Ma volcanic unit (Huang et al., 2005). The ca. 800 Ma pole puts Tarim at a palaeolatitude comparable to the position as in the Rodinia Map, although the relative orientation is somewhat different. The younger pole places Tarim at a significantly lower palaeolatitude than that adopted in the Rodinia Map. As no fold test has been reported for either of the two poles, and both study regions are inside younger fold-and-thrust belts, we cannot rule out vertical-axis rotation which can only be identified through comparative study of similar-age rocks in other parts of Tarim.

2.6. North China

Based mainly on tectonostratigraphic correlations, Li et al. (1996) suggested that the North China craton was adjacent to Siberia in Rodinia. The connection may have started as early as ca. 1800 Ma, and did not break-up until after ca. 600 Ma. Reliable palaeomagnetic data at both ca. 1770 Ma (Halls et al., 2000) and ca. 1350 Ma (Wu et al., 2005) from North China lend support to such a configuration (Fig. 7b). However, data from younger Proterozoic rocks suggest a slightly different configuration, involving a ca. 90° vertical-axis rotation of the North China




Fig. 7. Position of the North China craton (a) in Rodinia (after Zhang et al., 2006), and (b) during Paleo- to Mesoproterozoic (after Wu et al., 2005). Note that palaeomagnetic data in (b) suggest that Laurentia, Siberia, Baltica and North China could have been together for the ca. 1800–1300 Ma interval (Wu et al., 2005).

craton in relation to Laurentia between ca. 1350 Ma and 1200 Ma (S. Zhang et al., 2006) (Fig. 7a). According to S. Zhang et al. (2006), North China and Laurentia could have shared a common path between ca. 1200 Ma and ca. 700 Ma, but split up by ca. 615 Ma. It is unclear whether it was Siberia or other continental blocks that filled the gap between the two cratons for either the 1800–1350 Ma or the 1200–700 Ma intervals. We have adopted the rotation parameters for the North China craton provided by S. Zhang et al. (2006) for the Rodinia Map (Figs. 7a and 8). The lack of Neoproterozoic plume record in North China (e.g., Lu et al., 2008b) is consistent with it being at distances from continents like Australia and South China in Rodinia.

2.7. Minor terranes

The fragmented nature of many minor continental terranes, as well as their usual lack of any palaeomagnetic constraint, made it more difficult to place these terranes in Rodinia with much confidence. However, it is important to consider these terranes because they could account for spaces and geological links between larger cratons in Rodinia.

Keppie and Ramos (1999) proposed that two Central American terranes – Oaxaquia (Mexico) and Chortis (Honduras and Guatemala) – were situated along the northern boundary of South America in their reconstruction for the Ediacaran (Vendian)–Cambrian boundary. Keppie and Ortega-Gutierrez (1999) suggested that these blocks originated as arcs in a Grenvillian ocean between Laurentia, Baltica, and Amazonia, and were caught in-between the colliding cratons. These blocks experienced high grade, collisional-style metamorphism during their terminal collisions among Amazonia, Laurentia, and Baltica by ca. 1000 Ma. We place the Oaxaquia and Chortis blocks along the northern margin of Amazonia, within the zone

of its collision with Baltica at ca. 1000 Ma. Our model is also constrained by palaeomagnetic data from Oaxaquia (Ballard et al., 1989).

The space between Siberia and Laurentia in Rodinia is filled with several continental blocks that rifted away from northern Laurentia during the Cretaceous opening of the Canadian basin. These include terranes in northern Alaska, northern Chukchi Peninsula, Wrangel Island, the New Siberian Islands, Severnaya Zemlya, northern Taimyr and the Chukchi Plateau (e.g., Zonenshain et al., 1990; Vernikovskiy, 1997; Bogdanov et al., 1998; Embry, 1998; Nokleberg et al., 2000; Vernikovskiy and Vernikovskaya, 2001; Natal'in, 2004; Drachev, 2004). Our poor knowledge of these terranes makes geological correlation between northern Laurentia and Siberia more difficult. The position of the Kara Plate (exposed on Severnaya Zemlya in northern Taimyr) is debatable (Metelkin et al., 2005b). Although detrital zircon grains from the base of the Kara stratigraphy indicate a Baltican affinity (Pease et al., 2005, 2006), there are some geological arguments for a north Laurentian affinity (Zonenshain et al., 1990; Natal'in et al., 1999). Exact positions of these minor blocks in Rodinia are unclear due to the lack of palaeomagnetic data.

Previous reconstructions placed the Barentsia plate (exposed in Eastern Svalbard) against the southern part of east Greenland using their similarities in Neoproterozoic sedimentary successions (e.g., Winchester, 1988; Fairchild and Hambrey, 1995; Harland, 1997; Andresen, 2004). However, we choose to place it close to the northern part of East Greenland after Gee and Teben'kov (2004) who correlated the geochronological, structural, and stratigraphical data of Svalbard to northern Greenland. We have also reduced the size of the Barentsia plate in accordance with the recent discovery of a Caledonian arm in the Barents Sea (Breivik et al., 2002).




Fig. 8. A simplified (and reduced) Rodinia Map with legends.

3. Synthesis and animation: assembly and break-up of Rodinia, and formation of Gondwanaland

Through a series of cartoons, we illustrate a feasible scenario for the formation of Rodinia as shown in Fig. 8 and Appendix I, its break-up, and the eventual formation of Gondwanaland by the Early Cambrian. The rotation parameters for the major continental blocks are given in Appendix III, available online. A digital animation of the palaeogeographic evolution between 1100 Ma and 530 Ma, with geological features shown for selected time windows, is also given (Appendix II). We recognise that this is just one of a number of feasible scenarios for the evolution of Rodinia, and there are time intervals in the animation (e.g., the ca. 1000–820 Ma interval) for which we have very little palaeomagnetic constraints. Such reconstructions emphasise potential geody-

namic linkages and provide a testable hypothesis predicting continental positions and plate interactions, and enable palaeoclimatic modellers to simulate continent-ocean-atmosphere interactions.

3.1. The formation of Rodinia (ca. 1100–900 Ma)

At 1100 Ma (Fig. 9a), Laurentia, Siberia, North China, Cathaysia (part of present day South China) and perhaps Rio de la Plata were already together, and the Yangtze craton had begun its oblique collision with Laurentia (at southern Cathaysia; Greentree et al., 2006). However, all other continental blocks were still separated from Laurentia by oceans. The Australian craton, including the East Antarctica part of the Mawson craton, had amalgamated by then. King Island, where a 1287 ± 18 Ma metamorphic age was recently reported (Berry et al., 2005), Tas-

mania, and the South Tasman Rise (1119 ± 9 Ma quartz syenite, Fioretti et al., 2005), could have been close to where Yangtze and Laurentia were colliding, thus receiving Laurentia-sourced sediments (Berry et al., 2001).

By ca. 1050 Ma (Fig. 9b), Kalahari had probably collided with southern Laurentia (see Section 2.3). Continued collision of the Yangtze craton with western Laurentia may have caused the 1090–1030 Ma metamor-






Fig. 9. Cartoons showing the assembly and break-up of Rodinia, and the formation of Gondwanaland. (a) 1100 Ma; (b) 1050 Ma; (c) 1000 Ma; (d) 900 Ma; (e) 825 Ma; (f) 780 Ma; (g) 750 Ma; (h) 720 Ma; (i) 630 Ma; (j) 600 Ma; (k) 550 Ma; (l) 530 Ma.

(d) 900 Ma



(e) 825 Ma



(f) 780 Ma





Fig. 9. (Continued)


phism of the mafic sills in the Belt-Purcell Supergroup (Anderson and Davis, 1995). Convergent margins were developed between most continents, as the oceanic lithosphere between them was consumed during the assembly of Rodinia.

At ca. 1000 Ma (Fig. 9c), all but India, Australia–East Antarctica and Tarim had assembled to be joined with Laurentia, whereas the Yangtze craton was still suturing to Cathaysia (part of Laurentia). The transpressional movement between Greater India and Western Australia may explain the 1100–1000 Ma

(g) 750 Ma



(h) 720 Ma



(i) 630 Ma



Fig. 9. (Continued)


metamorphic ages reported from the Pinjarra Orogen (Bruguier et al., 1999; Fitzsimons, 2003).

By ca. 900 Ma all major known continental blocks had aggregated to form the Rodinia supercontinent (Figs. 8 and 9d, and Appendix I). Evidence for ca. 900 Ma orogenic events include

the ca. 920–880 Ma arc volcanics and ophiolite obduction in the eastern Sibao Orogen of South China (Li et al., 2005), 950–900 Ma arc volcanics along the northern margin of the Yangtze craton (Ling et al., 2003), and the 990–900 Ma high-grade metamorphic events in both the Eastern Ghats Belt of India

(j) 600 Ma

(Laurentia at high-latitude option)



(k) 550 Ma



(l) 530 Ma




Fig. 9. (Continued).

and the corresponding Rayner Province in East Antarctica (e.g., Mezger and Cosca, 1999; Boger et al., 2000; Fitzsimons, 2000; Kelly et al., 2002). This may also be the time when the Tarim craton joined Australia, as indicated by the development of the Aksu blueschist which predates ca. 800 Ma mafic dyke intru-

sions and have ^{40}Ar – ^{39}Ar cooling ages of 872–862 Ma (Zhang L.F., unpublished data, as quoted in Chen et al., 2004).

Stresses induced by the ca. 900 Ma event probably caused reactivation of older orogens within Rodinia. In the Mackenzie Mountains region of northwestern Laurentia, there is evidence

for an east-west compressional event (the Corn Creek Orogeny) sometime between 1033 Ma and 750 Ma (Thorkelson, 2000; Thorkelson et al., 2005). In the southern Capricorn Orogen inside the West Australian craton, the Edmundian Orogeny continued until ca. 900 Ma as shown by ^{40}Ar – ^{39}Ar ages (e.g., Occhipinti and Reddy, 2005). In the King Leopold Orogen of the North Australian craton, there was also the Yampi orogenic event (Tyler et al., 1998; Tyler and Griffin, 1990) with a possible ^{40}Ar – ^{39}Ar age of ca. 900 Ma (Bodorkos and Reddy, 2005).

3.2. Superplume events, continental rifting, and the prolonged break-up process of Rodinia (ca. 860–570 Ma)

Palaeomagnetic constraints for the time interval between 900 Ma and 830 Ma are very poor. Little geological record is available within Rodinia from this time interval, apart from a small number of 870–850 Ma intrusions such as those in South China and Africa (X.H. Li et al., 2003b; Johnson et al., 2005). Ca. 845 Ma and 870 Ma bimodal intrusions have also been reported from the Scandinavian Caledonides (Paulsson and Andreasson, 2002) and the Scottish promontory of Laurentia (Dalziel and Soper, 2001), both interpreted as representing the beginning of Rodinia break-up. Z.X. Li et al. (2003b) suggested that these intrusions could have been the first sign of a Rodinia superplume, i.e. local anatectic melts due to enhanced thermal gradients above ascending plume-heads.

Widespread plume activity did not occur until ca. 825 Ma, as shown by mafic dyke swarms, intra-continental mafic–ultramafic intrusions, and felsic intrusions (resulting from crustal melt or magma differentiation). However, such magmatism is commonly found in the polar end of Rodinia only (Li et al., 2004), including Australia (Zhao et al., 1994; Wingate et al., 1998), South China (Li et al., 1999; X.H. Li et al., 2003a; Z.X. Li et al., 2003b), Tarim (Zhang et al., 2006), India (Radhakrishna and Mathew, 1996), Kalahari (Frimmel et al., 2001), and the Arabian–Nubian terranes (Stein and Goldstein, 1996; Teklay et al., 2002). In places these intrusive rocks are unconformably overlain by similar-aged rift-related volcanoclastic successions, suggesting syn-magmatic doming (e.g., Li et al., 1999). Li et al. (1999) and Z.X. Li et al. (2003b) interpreted this widespread, largely synchronous magmatic event as the first major episode of superplume events which eventually led to the break-up of Rodinia (Fig. 9e). They regard the globally common sedimentary hiatus between ca. 900–880 Ma and 820 Ma as partly due to plume-induced crustal unroofing.

Away from the suggested polar superplume, similar-aged gabbro and monzonite intrusions are reported from the Scandinavian Caledonides, which are interpreted as indicative of continental rifting (Reginiussen et al., 1995). However, it is unclear whether this event was related to the high-latitude superplume event as no such rocks have been reported in-between the two regions. Nonetheless, such a connection would be possible given the mechanism for the generation of the superplume (see discussions below).

We illustrate the possible formation and tomography of a superplume (Fig. 10). Although thermal insulation of a supercontinent may elevate the temperature of the upper mantle (e.g.,

Anderson, 1982; Gurnis, 1988), Li et al. (2004) suggested that the dominant driving force for the generation of mantle superplumes (or superswells) is the double (circum-Rodinia) push-up effects of mantle avalanches (e.g., Kellogg et al., 1999; Moores et al., 2000; Fig. 10). Such avalanches could occur once oceanic slabs subducting below margins of the supercontinent become too dense to be supported at the mantle transitional zone (e.g., Tackley et al., 1993). On the surface the superplume would appear as a “plume cluster” (Ernst and Buchan, 2002; Schubert et al., 2004) consisting of “secondary” plumes originating above the top of the superplume (Courtillot et al., 2003).

Such a mechanism would explain the bipolar nature of superplume occurrences, as illustrated by the present-day Africa and Pacific superplumes (e.g., Zhao, 2001; Courtillot et al., 2003), which are likely the residuals of the Pangaeian superplume (e.g., Anderson, 1982; Burke and Torsvik, 2004) and the Palaeo-Pacific superplume (e.g., Larson, 1991a). The model would thus predict the occurrence of numerous oceanic plateaux and seamounts in the oceans on the opposite side of the Earth from Rodinia, traces of which would be found in the “Pan-African”-age orogens (e.g., parts of the Arabian–Nubian Shield; Stein and Goldstein, 1996). This model agrees with that of Condie (1998, 2000) in as much as superplumes are results of slab avalanches. It differs from the Condie (1998, 2000) model in that the superplume events in our model occurred after the formation of the supercontinent, not during the formation of the supercontinent. In our model the superplume events eventually led to the break-up of the supercontinent.

The ca. 825 Ma superplume event, followed by continental rifting, lingered on for ca. 25 million years, with another weaker magmatic peak occurring at ca. 800 Ma (Z.X. Li et al., 2003b; Ernst et al., 2008).

There appears to have been a global hiatus in plume/rift magmatism at around 790 ± 5 Ma (Z.X. Li et al., 2003b), but another superplume breakout occurred at ca. 780 Ma, with the best examples being the Gunbarrel event in Western Laurentia (Harlan et al., 2003b) and the Kangding event in South China (Z.X. Li et al., 2003b; Lin et al., 2007). By that time Rodinia had moved away from the northern polar region (Fig. 9f). India may have already rifted away from Rodinia if we accept that the palaeomagnetic pole from the Malani igneous suite (Torsvik et al., 2001b) is reliable and applicable for the entire 770–750 Ma interval, although no record of continental rifting and break-up has been reported for that time interval. The rapid rotation of Rodinia between ca. 820–800 Ma and ca. 780–750 Ma is interpreted by Li et al. (2004) as a true polar wander (TPW) event related to the occurrence of a high-latitude superplume (Evans, 1998, 2003b; Z.X. Li et al., 2003b). It is interesting to note that the rotation axis for the suggested TPW event is just east of Greenland (Li et al., 2004, Fig. 5), close to the Scandinavian Caledonides and the Scottish promontory of Laurentia where ca. 845 Ma and 870 Ma bimodal intrusions are present and are interpreted to represent the beginning of Rodinia break-up (Dalziel and Soper, 2001; Paulsson and Andreasson, 2002). Perhaps these magmatic events indicate the arrival of a plume-head which acted as the equatorial minimum-inertial axis for multiple TPW event(s) (Evans, 2003b; Li et al., 2004; Maloof et al., 2006).




Fig. 10. A cartoon showing the formation of a mantle “superplume” beneath supercontinent Rodinia (inspired by Kellogg et al. (1999) and Moores et al. (2000)).

Our proposed mechanism for the formation of the Rodinia superplume (i.e. due to mantle avalanches surrounding the supercontinent, possibly enhanced by the thermal insulation effect of the supercontinent; Fig. 10), and the palaeomagnetic implication that both the superplume and Rodinia above it may have moved from a high-latitude position to an equatorial position between ca. 820–800 Ma and ca. 780–750 Ma (Li et al., 2004), imply that whole-mantle convection is primarily driven by subduction processes (a kind of “top-down tectonics”, but differs from that of Anderson (2001) and others (e.g., Foulger et al., 2005) in that we have whole-mantle convection here), not by stationary heat sources at the core–mantle boundary.

By ca. 750 Ma (Fig. 9g), the western half of Rodinia may have started to break apart above an equatorial superplume. Bimodal magmatism at ca. 755–750 Ma was the last such major event seen across Rodinia (or a breaking-apart Rodinia). Like the ca. 825 Ma episode, syn-magmatic continental unroofing, documented by erosional contacts between ca. 770–750 Ma intrusive rocks and ca. 750–740 Ma rift successions, are common features (e.g., in western Kalahari, South China, and western and southeastern margins of Laurentia; see Z.X. Li et al., 2003b and references therein). By ca. 720 Ma (Fig. 9h), Australia–East Antarctica and South China are probably separated from each other by wide oceans. Even Kalahari and Siberia may have started to break away from Laurentia by this time. The ca. 750–700 Ma interval is also when the first major global Sturtian glaciation occurred and when most continents

were located at low- to moderate-latitude positions (the first snowball-Earth? Kirschvink, 1992; Hoffman et al., 1998). At ca. 650–630 Ma, when the dispersing continental blocks became even further aligned along the palaeo-equator (Fig. 9i), the second widespread low-latitude glaciation (the Marinoan glaciation, and another snowball-Earth event? Hoffman and Schrag, 2002) occurred. Numerous potential causes for these enigmatic global climatic events have been proposed and all are highly controversial. We refer readers to a dedicated web site (<http://snowballearth.org>) on current understanding of the subject, and the bibliography listed therein.

By ca. 600 Ma, the Amazonia, West Africa and Congo–São Francisco cratons had largely come together during the Brasiliano Orogeny (e.g., Trompette, 1997) (but note the alternative suggestion of Trindade et al., 2006 that these did not collide until Cambrian times by a Pampean–Araguaia orogeny), while Amazonia and Rio de la Plata were probably still attached to Laurentia (Fig. 9j). Siberia (Pisarevsky et al., 2008), North China (Zhang et al., 2006) and Baltica (Cawood and Pisarevsky, 2006; Pease et al., 2005, 2006) were separated from Laurentia by ca. 600 Ma. However, equivocal 580–560 Ma palaeomagnetic data put Laurentia at either a high-latitude position, or a low-latitude position (Cawood and Pisarevsky, 2006 and references therein). In our model we show the high-latitude option as the main figure, but have the low-latitude alternative shown as an insert (Fig. 9j). The high-latitude option makes it easier to explain the ca. 580 Ma Gaskiers glaciation in Baltica (e.g., Meert and van der

Voo, 1994), but this does not apply to the likely post-Marinoan glaciation in Tarim (Xiao et al., 2004) which lacks precise palaeomagnetic constraints. Amazonia was likely separated from Laurentia by ca. 570 Ma (e.g., Cawood and Pisarevsky, 2006 and references therein). The rifting away of continental blocks (e.g., Siberia, Kalahari, Rio de la Plata, Amazonia and Baltica) from Laurentia between ca. 630 Ma and 550 Ma may have been responsible for the widespread subsidence recorded at the margins of many of these cratons (e.g., Bond et al., 1984).

3.3. The birth of Gondwanaland (600–530 Ma)

Hoffman (1991) first suggested that the break-up of the Rodinia supercontinent involved fragmentation around Laurentia with continental pieces moving away from Laurentia and colliding on the other side of the Earth to form Gondwanaland. As mentioned in Section 3.2, with the exception of Kalahari and perhaps some minor terranes, West Gondwana was largely together by ca. 600 Ma. However, oceans still existed between Australia–East Antarctica, India, eastern Africa and Kalahari at that time (e.g., Meert, 2003; Jacobs and Thomas, 2004; Collins and Pisarevsky, 2005). By ca. 550 Ma (Fig. 9k), India had moved closer to its Gondwanaland position along the western margin of Australia, as recorded by the sinistral strike-slip movement along the Pinjarra Orogen (Fitzsimons, 2003). Kalahari started to collide with Congo and Rio de la Plata, thus closing the Neoproterozoic Adamastor Ocean between them (e.g., Prave, 1996). North China, separating from Laurentia–Siberia after ca. 650 Ma (S. Zhang et al., 2006), was drifting across the palaeo-Pacific Ocean toward Australia at that time.

Gondwanaland finally amalgamated by ca. 540–530 Ma (Fig. 9l) through the closure of both the so-called “Mozambique Ocean”, causing the Malagasy Orogeny in the East African Orogen (Meert, 2003; Jacobs and Thomas, 2004; Collins and Pisarevsky, 2005), and the final docking of India to Australia–East Antarctica along the Pinjarra Orogen (Fitzsimons, 2003; Boger and Miller, 2004; Collins and Pisarevsky, 2005; also known as the Kuunga Orogen, Meert and Van der Voo, 1996; Meert, 2003). The formation of Gondwanaland by ca. 530 Ma is supported by palaeomagnetic analyses (e.g., Meert and Van der Voo, 1996; Li and Powell, 2001). Both the South China and North China blocks were located close to Australia, as indicated by episodic bioprovince connections with east Gondwana during the Early to mid-Palaeozoic (e.g., Burrett and Richardson, 1980).

4. Concluding remarks

The late Mesoproterozoic and Neoproterozoic period is one of the most remarkable time intervals in Earth’s history. During this time we see the assembly and break-up of the supercontinent Rodinia that was ancestral to the long-lived supercontinent of Gondwanaland, possible global superplume events and rapid true polar wander event(s), repeated low-latitude glaciations, and finally the explosion of multicellular life (McMenamin and McMenamin, 1990) and the emergence of a plate dynamic and

climatic system very similar to what we have today (Moores, 2002; Evans, 2003a). The occurrence of some of these events, and possible genetic links between them, are still highly controversial. Nonetheless, these conceptual ideas enable us to think beyond the Phanerozoic “comfort zone” that persisted until recently. They have also forced geoscientists to think beyond their specific disciplines, bringing together tectonicists, structural geologists, geochemists, stratigraphers, petrologists, geophysicists, climatologists, and palaeontologists alike to work together toward a better understanding of Earth system science at the dawn of the Phanerozoic. With this background in mind, we present this overview of the formation and break-up of Rodinia. The scenarios presented here are not necessarily correct and there are many other scenarios which may explain some of the observations as well. It nonetheless provides a basis for future research on related topics.

Rodinia and Pangaea are the two supercontinents that we know to have included almost all the continents on Earth. Despite the secular nature of Earth’s evolution (and thus its tectonic processes), there are remarkable similarities between them: (1) They both had a lifespan of ca. 150 Ma: ca. 900 Ma to ≥ 750 Ma for Rodinia (this work), and ca. 320 Ma to 180–160 Ma for Pangaea (e.g., Li and Powell, 2001; Veevers, 2004); (2) The break-up of both supercontinents started with broad mantle upwellings (or superplumes) beneath them, resulting in widespread bimodal magmatism (including plume magmatism) and continental rifting (see Section 3.2); (3) The formation and break-up of Pangea also coincided with the occurrence of geomagnetic superchrons (Larson, 1991b; Eide and Torsvik, 1996), although there are still inadequate data for assessing the presence of similar events during the history of Rodinia. The Rodinia and Pangea supercontinental episodes thus provide us rare opportunities to gain insights into the 4D geodynamic system featuring close interactions between the thermal dynamics of the Earth’s outer core, mantle convections, and lithospheric plate tectonics.

Acknowledgments

Not all co-authors agree on everything written in this paper, and the map of Rodinia presented in this paper reflects only one of the plausible models. We are grateful for support from UNESCO/IUGS/IGCP and their national committees, the Australian Research Council (through the Tectonics Special Research Centre and Discovery Project grant DP0450020), the National Natural Science Foundation of China and the Swedish Research Council. This work would not have been possible without the inspiration and leaderships of the late professors Chris Powell, Raphael Unrug, and Henri Kampunzu. We also acknowledge the contributions made by all members of IGCP 440, particularly J. Myers, M.K. Pandit, J.P. Milesi, Guojie Hao and Guanghua Niu who helped to compile the individual map sheets, and other contributors as shown on the Rodinia Map (Appendix I). The palaeogeographic reconstructions were made using PLATES developed at the University of Texas at Austin. Tectonics Special Research Centre publication #387, and a contribution to IGCP 440.

Appendix I

The Geodynamic Map of Rodinia – see printed version inside the back cover of this volume; an electronic version is available at doi:10.1016/j.precamres.2007.04.021.

Appendix II

From Rodinia to Gondwanaland – An animated history, 1100–530 Ma – PowerPoint animation available at (link to the PowerPoint file at the Elsevier online version) doi:10.1016/j.precamres.2007.04.021.

Appendix III

Rotation parameters for selected time slices during the assembly and break-up of Rodinia and the assembly of Gondwanaland – available at doi:10.1016/j.precamres.2007.04.021.

References

- Åhäll, K.-I., Connely, J., 1998. Intermittent 1.53–1.13 Ga magmatism in western Baltica; age constraints and correlations within a postulated supercontinent. *Precambrian Res.* 92, 1–20.
- Aleinikoff, J.N., Zartman, R.E., Walter, M., Rankin, D.W., Lyttle, P.T., Burton, W.C., 1995. U–Pb ages of metarhyolites of the Catocin and Mount Rogers formations Central and Southern Appalachians; evidence for two pulses of Iapetan rifting. *Am. J. Sci.* 295, 428–454.
- Anderson, D.L., 1982. Hotspots, polar wander, Mesozoic convection and the geoid. *Nature* 297, 391–393.
- Anderson, D.L., 2001. Top-down tectonics? *Nature* 293, 2016–2018.
- Anderson, H.E., Davis, D.W., 1995. U–Pb geochronology of the Moyie sills, Purcell Supergroup, southeastern British Columbia: implications for the Mesoproterozoic geological history of the Purcell (Belt) basin. *Can. J. Earth Sci.* 32, 1180–1193.
- Andresen, A., 2004. On the origin of the Caledonian terranes of Svalbard. *Geologiska Föreningens Förhandlingar Abstract* 126, 77.
- Ashwal, L.D., Demaiffe, D., Torsvik, T.H., 2002. Petrogenesis of Neoproterozoic granitoids and related rocks from the Seychelles: the case for an Andean-type arc origin. *J. Petrol.* 43, 45–83.
- Ballard, M.M., Van der Voo, R., Urrutia-Fucugauchi, J., 1989. Paleomagnetic results from Grenvillian-aged rocks from Oaxaca Mexico: evidence for a displaced terrane. *Precambrian Res.* 42, 343–352.
- Bell, R.T., Jefferson, C.W., 1987. An hypothesis for an Australia–Canadian connection in the Late Proterozoic and the birth of the Pacific Ocean. *Proc. Pacific Rim Congress* 87, 39–50.
- Berry, R.F., Jenner, G.A., Meffre, S., Tubrett, M.N., 2001. A North American provenance for Neoproterozoic to Cambrian sandstones in Tasmania? *Earth Planet. Sci. Lett.* 192, 207–222.
- Berry, R.F., Holm, O.H., Steele, D.A., 2005. Chemical U–Th–Pb monazite dating and the Proterozoic history of King Island, southeast Australia. *Aust. J. Earth Sci.* 52, 461–471.
- Betts, P.G., Giles, D., 2006. The 1800–1100 Ma tectonic evolution of Australia. *Precambrian Res.* 144, 92–125.
- Bleeker, W., Ernst, R.E., 2006. Short-lived mantle generated magmatic events and their dyke swarms: the key unlocking Earth's paleogeographic record back to 2.6 Ga. In: E. Hanks, S. Mertanen, T. Rämö, J. Vuollo (Eds.), *Dyke Swarms—Time Markers of Crustal Evolution*. Taylor and Francis/Balkema, London, pp. 3–26.
- Blewett, R.S., Black, L.P., Sun, S.S., Knutson, J., Hutton, L.J., Bain, J.H.C., 1998. U–Pb zircon and Sm–Nd geochronology of the Mesoproterozoic of North Queensland; implications for a Rodinian connection with the Belt Supergroup of North America. *Precambrian Res.* 89, 101–127.
- Bodorkos, S., Reddy, S.M., 2005. $^{40}\text{Ar}/^{39}\text{Ar}$ constraints on Proterozoic cooling and exhumation of the Halls Creek orogen, WA. In: Wingate, M.T.D., Pisarevsky, S.A. (Eds.), *Supercontinent and Earth Evolution Symposium*. Geol. Soc. Aus. Inc., Fremantle, p. 34, Abs. No. 81.
- Bogdanov, N.A., Khain, V.Ye., Rosen, O.M., Shipilov, E.V., Vernikovskiy, V.A., Drachev, S.S., Kostyuchenko, S.L., Kuzmichev, A.B., Sekretov, S.B., 1998. Explanatory Notes for the Tectonic Map of the Kara and Laptev Seas and Northern Siberia (scale 1: 2,500,000). *Inst. Lithosphere, Moscow*, p. 116.
- Bogdanova, S.V., Page, L.M., Skridlaite, G., Taran, L.N., 2001. Proterozoic tectonothermal history in the western part of the East European Craton: $^{40}\text{Ar}/^{39}\text{Ar}$ geochronological constraints. *Tectonophysics* 339 (1–2), 39–66.
- Bogdanova, S.V., Bingen, B., Gorbatschev, R., Kheraskova, T.N., Kozlov, V.I., Puchkov, V.N., Volozh, Yu.A., 2008. The Eastern European Craton (Baltica) before and during the assembly of Rodinia. *Precambrian Res.* 160, 23–45.
- Boger, S.D., Miller, J.M., 2004. Terminal suturing of Gondwana and the onset of the Ross–Delamerian Orogeny: the cause and effect of an Early Cambrian reconfiguration of plate motions. *Earth Planet. Sci. Lett.* 219, 35–48.
- Boger, S.D., Carson, C.J., Wilson, C.J.L., Fanning, C.M., 2000. Neoproterozoic deformation in the Radok Lake region of the northern Prince Charles Mountains East Antarctica; evidence for a single protracted orogenic event. *Precambrian Res.* 104, 1–24.
- Boger, S.D., Wilson, C.J.L., Fanning, C.M., 2001. Early Paleozoic tectonism within the East Antarctic Craton; the final suture between East and West Gondwana? *Geology* 29, 463–466.
- Bond, G.C., Nickeson, P.A., Kominz, M.A., 1984. Breakup of a supercontinent between 625 Ma and 555 Ma: new evidence and implications for continental histories. *Earth Planet. Sci. Lett.* 70, 325–345.
- Borg, S.G., DePaolo, D.J., 1994. Laurentia, Australia, and Antarctica as a Late Proterozoic supercontinent: constraints from isotopic mapping. *Geology* 22, 307–310.
- Breivik, A.J., Mjelde, R., Grogan, P., Shimamura, H., Murai, Y., Nishimura, Y., Kuwano, A., 2002. A possible Caledonide arm through the Barents Sea imaged by OBS data. *Tectonophysics* 355, 67–97.
- Briden, J.C., Duff, B.A., Kröner, A., 1979. Palaeomagnetism of the Koras Group, Northern Cape Province South Africa. *Precambrian Res.* 10, 43–57.
- Brookfield, M.E., 1993. Neoproterozoic Laurentia–Australia fit. *Geology* 21, 683–686.
- Brown, L.L., McEnroe, S.A., 2004. Palaeomagnetism of the Egersund–Ogna anorthosite, Rogaland, Norway, and the position of Fennoscandia in the late proterozoic. *Geophys. J. Int.* 158, 479–488.
- Bruguier, O., Bosch, D., Pidgeon, R.T., Byrne, D.I., Harris, L.B., 1999. U–Pb chronology of the Northampton Complex, Western Australia: evidence for Grenvillian sedimentation, metamorphism and deformation and geodynamic implications. *Contrib. Mineral. Petrol.* 136, 258–272.
- Buchan, K.L., Dunlop, D.J., 1976. Paleomagnetism of the Haliburton Intrusions: superimposed magnetisations, metamorphism, and tectonics in the late Precambrian. *J. Geophys. Res.* 81, 2951–2966.
- Buchan, K.L., Halls, H.C., 1990. Paleomagnetism of Proterozoic mafic dyke swarms of the Canadian Shield. In: Parker, A.J., Rickwood, P.C., Tucker, D.H. (Eds.), *Mafic Dykes and Emplacement Mechanisms*. Balkema, Rotterdam, pp. 209–230.
- Buchan, K.L., Mertanen, S., Park, R.G., Pesonen, L.J., Elming, S.A., Abrahamson, N., Bylund, G., 2000. Comparing the drift of Laurentia and Baltica in the Proterozoic: the importance of key palaeomagnetic poles. *Tectonophysics* 319, 167–198.
- Buchan, K.L., Ernst, R.E., Hamilton, M.A., Mertanen, S., Pesonen, L.J., Elming, S.A., 2001. Rodinia: the evidence from integrated palaeomagnetism and U–Pb geochronology. *Precambrian Res.* 110, 9–32.
- Burke, K., Torsvik, T.H., 2004. Derivation of Large Igneous Provinces of the past 200 million years from long-term heterogeneities in the deep mantle. *Earth Planet. Sci. Lett.* 227, 531–538.
- Burrett, C., Berry, R., 2000. Proterozoic Australia–Western United States (AUSWUS) fit between Laurentia and Australia. *Geology* 28, 103–106.
- Burrett, C., Richardson, R., 1980. Trilobite biogeography and Cambrian tectonic models. *Tectonophysics* 63, 155–192.
- Bylund, G., Elming, S.-Å., 1992. The Dala dolerites, central Sweden, and their palaeomagnetic signature. *GFF* 114, 143–153.

- Calver, C.R., Walter, M.R., 2000. The late Neoproterozoic Grassy Group of King Island Tasmania; correlation and palaeogeographic significance. *Precambrian Res.* 100, 299–312.
- Cawood, P.A., Pisarevsky, S.A., 2006. Was Baltica right-way-up or upside-down in the Neoproterozoic? *J. Geol. Soc. Lond.* 163, 753–759.
- Cawood, P.A., McCausland, P.J.A., Dunning, G.R., 2001. Opening Iapetus: constraints from the Laurentian margin in Newfoundland. *Geol. Soc. Am. Bull.* 113, 443–453.
- Čečys, A., Benn, K., 2007. Emplacement and deformation of the ca. 1.45 Ga Karlshamn granitoid pluton, southeastern Sweden, during ENE-WSW Danopolonian shortening. *Int. J. Earth Sci. (Geologische Rundschau)* 96, 397–414.
- Čečys, A., Bogdanova, S., Jansson, C., Bibikova, E., Kornfält, K.-A., 2002. The Stenshuvud and Tåghusa granitoids: new representatives of Mesoproterozoic magmatism in southern Sweden. *GFF* 124, 149–162.
- Chen, Y., Xu, B., Zhan, S., Li, Y.G., 2004. First mid-Neoproterozoic paleomagnetic results from the Tarim Basin (NW China) and their geodynamic implications. *Precambrian Res.* 133, 271–281.
- Collins, A.S., 2003. Structure and age of the Northern Leeuwin Complex Western Australia: constraints from field mapping and U–Pb isotopic analysis. *Aust. J. Earth Sci.* 50, 585–599.
- Collins, A.S., 2006. Madagascar and the amalgamation of Central Madagascar. *Gond. Res. (GR Focus)* 9, 3–16.
- Collins, A.S., Pisarevsky, S.A., 2005. Amalgamating eastern Gondwana: the evolution of the Circum-Indian Orogens. *Earth-Sci. Rev.* 71, 229–270.
- Collins, A.S., Windley, B.F., 2002. The tectonic evolution of central and northern Madagascar and its place in the final assembly of Gondwana. *J. Geol.* 110, 325–339.
- Collins, A.S., Fitzsimons, I.C.W., Hulscher, B., Razakamanana, T., 2003. Structure of the eastern margin of the East African Orogen in central Madagascar. *Precambrian Res.* 123, 111–133.
- Condie, K.C., 1998. Episodic continental growth and supercontinents; a mantle avalanche connection? *Earth Planet. Sci. Lett.* 163, 97–108.
- Condie, K.C., 2000. Episodic continental growth models: afterthoughts and extensions. *Tectonophysics* 322, 153–162.
- Condie, K.C., Rosen, O.M., 1994. Laurentia–Siberia connection revisited. *Geology* 22, 168–170.
- Courtillot, V., Davaille, A., Besse, J., Stock, J., 2003. Three distinct types of hotspots in the Earth's mantle. *Earth Planet. Sci. Lett.* 205, 295–308.
- Cox, R., Coleman, D.S., Chokel, C.B., DeOreo, S.B., Wooden, J.L., Collins, A.S., De Waele, B., Kröner, A., 2004. Proterozoic tectonostratigraphy and palaeogeography of central Madagascar derived from detrital zircon U–Pb age populations. *J. Geol.* 112, 379–399.
- D'Agrella-Filho, M.S., Pacca, I.G., Renne, P.R., Onstott, T.C., Teixeira, W., 1990. Paleomagnetism of Middle Proterozoic (1.01 to 1.08 Ga) mafic dykes in southeastern Bahia State–São Francisco Craton, Brazil. *Earth Planet. Sci. Lett.* 101, 332–348.
- D'Agrella-Filho, M.S., Pacca, I.I.G., Trindade, R.I.F., Teixeira, W., Raposo, M.I.B., Onstott, T.C., 2004. Paleomagnetism and Ar-40/Ar-39 ages of mafic dikes from Salvador (Brazil): new constraints on the São Francisco craton APW path between 1080 and 1010 Ma. *Precambrian Res.* 132, 55–77.
- Dalziel, I.W.D., 1991. Pacific margins of Laurentia and East Antarctica–Australia as a conjugate rift pair: evidence and implications for an Eocambrian supercontinent. *Geology* 19, 598–601.
- Dalziel, I.W.D., 1997. Neoproterozoic–Paleozoic geography and tectonics: review, hypothesis, environmental speculation. *Geol. Soc. Am. Bull.* 109, 16–42.
- Dalziel, I.W.D., Soper, N.J., 2001. Neoproterozoic extension on the Scottish Promontory of Laurentia; paleogeographic and tectonic implications. *J. Geol.* 109, 299–317.
- Davidson, A., 1995. A review of the Grenville orogen in its North American type area. *J. Aust. Geol. Geophys.* 16, 3–24.
- Davidson, A., 2008. Late paleoproterozoic to mid-neoproterozoic history of northern laurentia: An overview of central rodinia. *Precambrian Res.* 160, 5–22.
- Davis, D.W., Green, J.C., 1997. Geochronology of the North American Mid-continent rift in western Lake Superior and implications for its geodynamic evolution. *Can. J. Earth Sci.* 34, 476–488.
- Davis, D.W., Paces, J.B., 1990. Time resolution of geologic events on the Keweenaw Peninsula and applications for development of the Midcontinent Rift system. *Earth Planet. Sci. Lett.* 97, 54–64.
- Davis, D.W., Sutcliffe, R.H., 1985. U–Pb ages from the Nipigon plate and northern Lake Superior. *Geol. Soc. Am. Bull.* 96, 1572–1579.
- Dawson, G.C., Krapez, B., Fletcher, I.R., McNaughton, N.J., Rasmussen, B., 2003. 1.2 Ga thermal metamorphism in the Albany–Fraser Orogen of Western Australia: consequence of collision or regional heating by dyke swarms? *J. Geol. Soc. Lond.* 160, 29–37.
- De Waele, B., 2005. The Proterozoic geological history of the Irumide belt, Zambia. PhD Thesis, Curtin University of Technology, Perth, 468 pp.
- De Waele, B., Johnson, S.P., Pisarevsky, S.A., 2008. Palaeoproterozoic to Neoproterozoic growth and evolution of the Eastern Congo Craton: Its role in the Rodinia puzzle. *Precambrian Res.* 160, 127–141.
- De Waele, B., Kampunzu, A.B., Mapani, B.S.E., Tembo, F., 2006a. The Irumide Belt of Zambia. *J. Afr. Earth Sci.* 46, 36–70.
- De Waele, B., Liégeois, J.P., Nemchin, A.A., Tembo, F., 2006b. Isotopic and geochemical evidence of Proterozoic episodic crustal reworking within the Irumide Belt of south-central Africa, the southern metacratonic boundary of an Archaean Bangweulu Craton. *Precambrian Res.* 148, 225–256.
- De Waele, B., Wingate, M.T.D., Fitzsimons, I.C.W., Mapani, B.S.E., 2003. Untying the Kibaran knot; a reassessment of Mesoproterozoic correlations in Southern Africa based on SHRIMP U–Pb data from the Irumide Belt. *Geology* 31, 509–512.
- Deblond, A., Punzalan, L.E., Boven, A., Tack, L., 2001. The Malagarazi Supergroup of southeast Burundi and its correlative Bukoba Supergroup of northwest Tanzania: Neo- and Mesoproterozoic chronostratigraphic constraints from Ar–Ar ages on mafic intrusive rocks. *J. Afr. Earth Sci.* 32, 435–449.
- Diehl, J.F., Haig, T.D., 1994. A paleomagnetic study of the lava flows within the Copper Harbour Conglomerate Michigan: new results and implications. *Can. J. Earth Sci.* 31, 369–380.
- Direen, N.G., Crawford, A.J., 2003. The Tasman Line: where is it, what is it, and is it Australia's Rodinian breakup boundary? *Aust. J. Earth Sci.* 50, 491–502.
- Drachev, S.S., 2004. Siberian Arctic continental margin: constraints and uncertainties of plate tectonic model. In: AGU Fall Meet. Suppl., (Abstract GP43C-07).
- Eglington, B.M., Armstrong, R.A., 2003. Geochronological and isotopic constraints on the Mesoproterozoic Namaqua–Natal Belt: evidence from deep borehole intersections in South Africa. *Precambrian Res.* 125, 179–189.
- Eide, E.A., Torsvik, T.H., 1996. Paleozoic supercontinental assembly, mantle flushing, and genesis of the Kiaman Superchron. *Earth Planet. Sci. Lett.* 144, 389–402.
- Eisbacher, G.H., 1985. Late Proterozoic rifting, glacial sedimentation, and sedimentary cycles in the light of Windermere deposition, western Canada. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 51, 231–254.
- Elming, S.A., Pesonen, L.J., Leino, M.A.H., Khranov, A.N., Mikhailova, N.P., Krasnova, A.F., Mertanen, S., Bylund, G., Terho, M., 1993. The Drift of the Fennoscandian and Ukrainian Shields during the Precambrian—a paleomagnetic analysis. *Tectonophysics* 223, 177–198.
- Elming, S.-Å., Mikhailova, N.P., Kravchenko, S., 2001. Palaeomagnetism of Proterozoic rocks from the Ukrainian Shield: new tectonic reconstructions of the Ukrainian and the Fennoscandian Shields. *Tectonophysics* 339, 19–38.
- Embleton, B.J.J., Williams, G.E., 1986. Low paleolatitude of deposition for Late Precambrian periglacial varvites in South Australia: implications for paleoclimatology. *Earth Planet. Sci. Lett.* 79, 419–430.
- Embry, A.F., 1998. Counterclockwise rotation of the Arctic Alaska plate: best available model or untenable hypothesis for the opening of the Amerasia Basin. *Polarforschung* 68, 247–255.
- Ernst, R.E., Buchan, K.L., 1993. Paleomagnetism of the Abitibi dike swarm, southern Superior Province, and implications for the Logan Loop. *Can. J. Earth Sci.* 30, 1886–1897.
- Ernst, R.E., Buchan, K.L., 2002. Maximum size and distribution in time and space of mantle plumes; evidence from large igneous provinces. In: Condie Kent, C., Abbott Dallas, H., Des Marais David, J. (Eds.), *Superplume Events in Earth History: Causes and Effects*. Pergamon Press, Oxford, United Kingdom.

- Ernst, R.E., Head, J.W., Parfitt, E., Grosfils, E., Wilson, L., 1995. Giant radiating dyke swarms on Earth and Venus. *Earth-Sci. Rev.* 39, 1–58.
- Ernst, R.E., Buchan, K.L., Hamilton, M.A., Okrugin, A.V., D., T.M., 2000. Integrated paleomagnetism and U–Pb geochronology of mafic dikes of the Eastern Anabar Shield Region Siberia: implications for mesoproterozoic paleolatitude of Siberia and comparison with Laurentia. *J. Geol.* 108, 381–401.
- Ernst, R.E., Wingate, M.T.D., Buchan, K.L., Li, Z.X., 2008. Global record of 1600–700 Ma Large Igneous Provinces (LIPs): Implications for the reconstruction of the proposed Nuna (Columbia) and Rodinia supercontinents. *Precambrian Res.* 160, 159–178.
- Evans, D.A.D., 1998. Supercontinental cycles and true polar wander. In: Bird, Robert, T., Powell, Christopher, McA., Wingate (Eds.), *The Assembly and Breakup of Rodinia; Proceedings of a Workshop. Abstracts. Geological Society of Australia, Sydney, NSW, Australia*, pp. 34–35.
- Evans, D.A.D., 2000. Stratigraphic, geochronological, and paleomagnetic constraints upon the Neoproterozoic climatic paradox. *Am. J. Sci.* 300, 347–433.
- Evans, D.A.D., 2003a. A fundamental Precambrian–Phanerozoic shift in earth's glacial style? *Tectonophysics* 375, 353–385.
- Evans, D.A.D., 2003b. True polar wander and supercontinents. *Tectonophysics* 362, 303–320.
- Evans, D.A.D., 2006. Proterozoic low orbital obliquity and axial-dipolar geomagnetic field. *Nature* 444, 51–55.
- Evans, D.A.D., Li, Z.X., Kirschvink, J.L., Wingate, M.T.D., 2000. A high-quality mid-Neoproterozoic paleomagnetic pole from South China, with implications for ice ages and the breakup configuration of Rodinia. *Precambrian Res.* 100, 313–334.
- Fairchild, I.J., Hambrey, M.J., 1995. Vendian basin evolution in East Greenland and NE Svalbard. *Precambrian Res.* 73, 217–233.
- Fanning, C.M., Link, P.K., 2004. U–Pb SHRIMP ages of Neoproterozoic (Sturtian) glaciogenic Pocahontas Formation, southeastern Idaho. *Geology* 32, 881–884.
- Fetter, A.H., Goldberg, S.A., 1995. Age and geochemical characteristics of bimodal magmatism in the Neoproterozoic Grandfather Mountain rift basin. *J. Geol.* 103, 313–326.
- Fioretti, A.M., Black, L.P., Foden, J., Visona, D., 2005. Grenville-age magmatism at the South Tasman Rise (Australia): a new piercing point for the reconstruction of Rodinia. *Geology* 33, 769–772.
- Fitzsimons, I.C.W., 2000. Grenville-age basement provinces in East Antarctica: evidence for three separate collisional orogens. *Geology* 28, 879–882.
- Fitzsimons, I.C.W., 2003. Proterozoic basement provinces of southern and southwestern Australia, and their correlation with Antarctica. In: Yoshida, M., Windley, B.F., Dasgupta, S. (Eds.), *Proterozoic East Gondwana: Supercontinent Assembly and Breakup. Special Publications, Geological Society, London*, pp. 93–130.
- Fitzsimons, I.C.W., Hulscher, B., 2005. Out of Africa: detrital zircon provenance of central Madagascar and Neoproterozoic terrane transfer across the Mozambique Ocean. *Terra Nova* 17, 224–235.
- Foulger, G.R., Natland, J.H., Presnall, D.C., Anderson, D.L. (Eds.), 2005. *Plates, Plumes and Paradigms. Geological Society of America Special Paper* 388, p. 881.
- Frimmel, H.E., Kloetzli, U.S., Siegfried, P.R., 1996. New Pb–Pb single zircon age constraints on the timing of Neoproterozoic glaciation and continental break-up in Namibia. *J. Geol.* 104, 459–469.
- Frimmel, H.E., Zartman, R.E., Spath, A., 2001. The Richtersveld Igneous Complex South Africa: U–Pb zircon and geochemical evidence for the beginning of Neoproterozoic continental breakup. *J. Geol.* 109, 493–508.
- Frost, B.R., Avchenko, O.V., Chamberlain, K.R., Frost, C.D., 1998. Evidence for extensive Proterozoic remobilization of the Aldan shield and implications for Proterozoic plate tectonic reconstructions of Siberia and Laurentia. *Precambrian Res.* 89, 1–23.
- Fuck, R.A., Brito Neves, B.B., Schobbenhaus, C., 2008. Rodinia descendants in South America. *Precambrian Res.* 160, 108–126.
- Gallet, Y., Pavlov, V.E., Semikhatov, M.A., Petrov, P.Yu., 2000. Late Mesoproterozoic magnetostratigraphic results from Siberia: paleogeographic implications and magnetic field behaviour. *J. Geophys. Res.* 105, 16481–16499.
- Gee, D.G., Teben'kov, A.M., 2004. Svalbard: a fragment of the Laurentian margin. In: Gee, D.G., Pease, V.L. (Eds.), *The Neoproterozoic Timanide Orogen of Eastern Baltica. Geol. Soc. London. Memoirs* 30, pp. 191–206.
- Gladkochub, D.P., Wingate, M.T.D., Pisarevsky, S.A., Donskaya, T.V., Mazukabzov, A.M., Ponomarchuk, V.A., Stanevich, A.M., 2006. Mafic intrusions in southwestern Siberia and implications for a Neoproterozoic connection with Laurentia. *Precambrian Res.* 147, 260–278.
- Glass, L.M., Phillips, D., 2006. The Kalkarindji continental flood basalt province: A new Cambrian large igneous province in Australia with possible links to faunal extinctions. *Geology* 34, 461–464.
- Goodge, J.W., Myrow, P., Williams, I.S., Bowring, S.A., 2002. Age and provenance of the Beardmore Group Antarctica: constraints on Rodinia supercontinent breakup. *J. Geol.* 110, 393–406.
- Gorbatshev, R., Bogdanova, S., 1993. Frontiers in the Baltic Shield. *Precambrian Res.* 64, 3–21.
- Gose, W.A., Helper, M.A., Connelly, J.N., Hutson, F.E., Dalziel, I.W.D., 1997. Paleomagnetic data and U–Pb isotopic age determinations from Coats Land Antarctica; implications for late Proterozoic plate reconstructions. *J. Geophys. Res.* B 102, 7887–7902.
- Gower, C.F., Rivers, T., Ryan, A.B., 1990. Mid-Proterozoic Laurentia–Baltica: an overview of its geological evolution and a summary of the contributions made by this volume. In: Gower, C.F., Rivers, T., Ryan, A.B. (Eds.), *Mid-Proterozoic Laurentia–Baltica. Geological Association of Canada*, pp. 1–22.
- Greentree, M.R., Li, Z.X., Li, X.H., Wu, H., 2006. Late Mesoproterozoic to earliest Neoproterozoic basin record of the Sibao orogenesis in western South China and relationship to the assembly of Rodinia. *Precambrian Res.* 151, 79–100.
- Grey, K., Corkeron, M., 1998. Late Neoproterozoic stromatolites in glaciogenic successions of the Kimberley region Western Australia: evidence for a second Marinoan glaciation. *Precambrian Res.* 92, 65–87.
- Groenewald, P.B., Moyes, A.B., Grantham, G.H., Krynauw, J.R., 1995. East Antarctic crustal evolution: geological constraints and modelling in Western Dronning Maud Land. *Precambrian Res.* 75, 231–250.
- Guo, Z.J., Yin, A., Robinson, A., Jia, C.Z., 2005. Geochronology and geochemistry of deep-drill-core samples from the basement of the central Tarim basin. *J. Asian Earth Sci.* 25, 45–56.
- Gurnis, M., 1988. Large-scale mantle convection and the aggregation and dispersal of supercontinents. *Nature* 332, 695–699.
- Halls, H.C., Pesonen, L.J., 1982. Paleomagnetism of Keweenaw rocks. *Geol. Soc. Am. Mem.* 156, 173–201.
- Halls, H.C., Li, J.H., Davis, D., Hou, G., Zhang, B.X., Qian, X.L., 2000. A precisely dated Proterozoic palaeomagnetic pole from the North China craton, and its relevance to palaeocontinental reconstruction. *Geophys. J. Int.* 143, 185–203.
- Halverson, G.P., Hoffman, P.F., Schrag, D.P., Maloof, A.C., Rice, A.H.N., 2005. Toward a Neoproterozoic composite carbon-isotope record. *Geol. Soc. Am. Bull.* 117, 1181–1207.
- Handke, M.J., Tucker, R.D., Ashwal, L.D., 1999. Neoproterozoic continental arc magmatism in west-central Madagascar. *Geology* 27, 351–354.
- Hanley, L.M., Wingate, M.T.D., 2000. SHRIMP zircon age for an Early Cambrian dolerite dyke: an intrusive phase of the Antrim Plateau Volcanics of northern Australia. *Aust. J. Earth Sci.* 47, 1029–1040.
- Hanson, R.E., Martin, M.W., Bowring, S.A., Munyanyiwa, H., 1998. U–Pb zircon age for the Umkondo dolerites, eastern Zimbabwe: 1.1 Ga large igneous province in southern Africa East Antarctica and possible Rodinia correlations. *Geology* 26, 1143–1146.
- Hanson, R.E., Crowley, J.L., Bowring, S.A., Ramezani, J., Gose, W.A., Dalziel, I.W.D., Pancake, J.A., Seidel, E.K., Blenkinsop, T.G., Mukwakwami, J., 2004. Coeval large-scale magmatism in the Kalahari and Laurentian cratons during Rodinia assembly. *Science* 304, 1126–1129.
- Hargrove, U.S., Stern, R.J., Kimura, J.I., Johnson, P.R., 2004. What is the significance of Adakitic Granitoids and Zircon inheritance in Juvenile Arc Rocks of the Neoproterozoic Makkah Batholith, Saudi Arabia? In: *AGU Fall Meet. Suppl., Eos Trans., San Francisco (Abstract V13B-1472)*.
- Harlan, S.S., Geissman, J.W., Snee, L.W., 1997. Paleomagnetic and $^{40}\text{Ar}/^{39}\text{Ar}$ geochronologic data from late Proterozoic mafic dykes and sills, Montana and Wyoming. *USGS Professional Paper* 1580, 16 pp.

- Harlan, S.S., Geissman, J.W., Premo, W.R., 2003a. Paleomagnetism and geochronology of an Early Proterozoic quartz diorite in the southern Rind River Range, Wyoming USA. *Tectonophysics* 362, 105–122.
- Harlan, S.S., Heaman, L., LeCheminant, A.N., Premo, W.R., 2003b. Gunbarrel mafic magmatic event: a key 780 Ma time marker for Rodinia plate reconstructions. *Geology* 31, 1053–1056.
- Harland, W.B., 1997. The geology of Svalbard. *Geol. Soc. Lond. Memoir*, 17.
- Heaman, L.M., 1997. Global mafic magmatism at 2.45 Ga: remnants of an ancient large igneous province? *Geology* 25, 299–302.
- Heaman, L.M., LeCheminant, A.N., Rainbird, R.H., 1992. Nature and timing of Franklin igneous events Canada; implications for a late Proterozoic mantle plume and the break-up of Laurentia. *Earth Planet. Sci. Lett.* 109, 117–131.
- Henry, S.G., Mauk, F.J., Van der Voo, R., 1977. Paleomagnetism of the upper Keweenaw sediments: the Nonesuch Shale and Freda Sandstone. *Can. J. Earth Sci.* 14, 1128–1138.
- Hodych, J.P., Cox, R.A., Košler, J., 2004. An equatorial Laurentia at 550 Ma confirmed by Grenvillian inherited zircons dated by LAM ICP-MS in the Skinner Cove volcanics of western Newfoundland: implications for inertial interchange true polar wander. *Precambrian Res.* 129, 93–113.
- Hoffman, P.F., 1991. Did the breakout of Laurentia turn Gondwanaland inside-out? *Science* 252, 1409–1412.
- Hoffman, P.F., Schrag, D.P., 2002. The snowball Earth hypothesis: testing the limits of global change. *Terra Nova* 14, 129–155.
- Hoffman, P.F., Hawkins, D.P., Isachsen, C.E., Bowring, S.A., 1996. Precise U–Pb zircon age for early Damaran magmatism in the Summas Mountains and Welwitschia inlier, northern Damara belt Namibia. *Comm. Geol. Surv. Namibia* 11, 47–52.
- Hoffman, P.F., Kaufman, A.J., Halverson, G.P., Schrag, D.P., 1998. A Neoproterozoic snowball earth. *Science* 281, 1342–1346.
- Hoffmann, K.H., Condon, D.J., Bowring, S.A., Crowley, J.L., 2004. U–Pb zircon date from the Neoproterozoic Ghaub Formation Namibia: constraints on Marinoan glaciation. *Geology* 32, 817–820.
- Holm, O.H., Crawford, A.J., Berry, R.F., 2003. Geochemistry and tectonic settings of meta-igneous rocks in the Arthur Lineament and surrounding area, northwest Tasmania. *Aust. J. Earth Sci.* 50, 903–918.
- Huang, B.C., Xu, B., Zhang, C.X., Li, Y.A., Zhu, R.X., 2005. Paleomagnetism of the Baiyisi volcanic rocks (ca. 740 Ma) of Tarim Northwest China: a continental fragment of neoproterozoic Western Australia? *Precambrian Res.* 142, 83–92.
- Hutton, L., Fanning, C.M., Garrad, P., 1996. Grenvillian age magmatic and metamorphic event in the Cape River area, North Queensland; significance for late Mesoproterozoic continental reconstructions. In: Kennard, John (Eds.), *Geoscience for the Community; 13th Australian Geological Convention*, *Geol. Soc. Aus. Abst. Geological Society of Australia*, Sydney, NSW, Australia, p. 209.
- Hyodo, H., York, D., Dunlop, D.J., 1993. Tectonothermal history in the Mattawa Area, Ontario, Canada deduced from paleomagnetism and Ar-40/Ar-39 dating of a Grenville Dike. *J. Geophys. Res.* 98 (B10), 18001–18010.
- Idnurm, M., Giddings, J.W., 1988. Australian Precambrian polar wander: a review. *Precambrian Res.* 40, 61–88.
- Idnurm, M., Giddings, J.W., 1995. Paleoproterozoic Neoproterozoic North-America Australia Link—new evidence from Paleomagnetism. *Geology* 23, 149–152.
- Jacobs, J., Thomas, R.J., 2004. Himalayan-type indenter-escape tectonics model for the southern part of the late Neoproterozoic-early Paleozoic East African–Antarctic orogen. *Geology* 32, 721–724.
- Jacobs, J., Bauer, W., Fanning, C.M., 2003a. New age constraints for Grenville-age metamorphism in western central Dronning Maud Land (East Antarctica), and implications for the palaeogeography of Kalahari in Rodinia. *Int. J. Earth Sci.* 92, 301–315.
- Jacobs, J., Fanning, C.M., Bauer, W., 2003b. Timing of Grenville-age vs Pan-African medium- to high grade metamorphism in western Dronning Maud Land (East Antarctica) and significance for correlations in Rodinia and Gondwana. *Precambrian Res.* 125, 1–20.
- Jacobs, J., Pisarevsky, S., Thomas, R.J., Becker, T., 2008. The Kalahari Craton during the assembly and dispersal of Rodinia. *Precambrian Res.* 160, 142–158.
- Jefferson, C.W., Parrish, R.R., 1989. Late Proterozoic stratigraphy, U–Pb zircon ages, and rift tectonics, Mackenzie Mountains, northwestern Canada. *Can. J. Earth Sci.* 26, 1784–1801.
- Jiang, G., Sohl, L.E., Christie-Blick, N., 2003. Neoproterozoic stratigraphic comparison of the Lesser Himalaya (India) and Yangtze block (south China): paleogeographic implications. *Geology* 31, 917–920.
- Johnson, S.P., De Waele, B., Liyungu, A.K., 2006. U–Pb SHRIMP geochronology of granitoid rocks in eastern Zambia: terrane subdivision of the Mesoproterozoic Southern Irumide Belt. *Tectonics* 25, TC6004, doi:10.1029/2006TC001777.
- Johnson, S.P., De Waele, B., Evans, D., Banda, W., Tembo, F., Milton, J.A., Tani, K., 2007a. Geochronology of the Zambezi Supracrustal sequence, southern Zambia: A record of Neoproterozoic divergent processes along the southern margin of the Congo Craton. *J. Geol.* 115, 355–374.
- Johnson, S.P., De Waele, B., Tembo, F., Katongo, C., Tani, K., Chang, Q., Iizuka, T., Dunkley, D., 2007b. Geochemistry, geochronology and isotopic evolution of the Chewore-Rufunsa terrane, Southern Irumide Belt: A Mesoproterozoic, continental-margin-arc. *J. Petrol.* 48, 1411–1441.
- Johnson, S.P., Rivers, T., De Waele, B., 2005. A review of the Mesoproterozoic to early Palaeozoic magmatic and tectonothermal history of south-central Africa: implications for Rodinia and Gondwana. *J. Geol. Soc.* 162, 433–450.
- Jones, D.L., Bates, M.P., Li, Z.X., Corner, B., Hodgkinson, G., 2003. Palaeomagnetic results from the ca. 1130 Ma Borgmassivet intrusions in the Ahlmannryggen region of Dronning Maud Land, Antarctica, and tectonic implications. *Tectonophysics* 375, 247–260.
- Kamo, S.L., Gower, C.F., 1994. Note: U–Pb baddeleyite dating clarifies age of characteristic paleomagnetic remanence of Long Range dykes, southeastern Labrador. *Atlantic Geol.* 30, 259–262.
- Karlstrom, K.E., Williams, M.L., McLelland, J., Geissman, J.W., Ahall, K.I., 1999. Refining Rodinia: geological evidence for the Australia–western U.S. connection in the Proterozoic. *GSA Today* 9, 1–7.
- Karlstrom, K.E., Bowring, S.A., Dehler, C.M., Knoll, A.H., Porter, S.M., Des Marais, D.J., Weil, A.B., Sharp, Z.D., Geissman, J.W., Elrick, M.B., Timmons, J.M., Crossey, L.J., Davidek, K.L., 2000. Chuar Group of the Grand Canyon: record of breakup of Rodinia, associated change in the global carbon cycle, and ecosystem expansion by 740 Ma. *Geology* 28, 619–622.
- Karlstrom, K.E., Ahall, K.I., Harlan, S.S., Williams, M.L., McLelland, J., Geissman, J.W., 2001. Long-lived (1.8–1.0 Ga) convergent orogen in southern Laurentia, its extensions to Australia and Baltica, and implications for refining Rodinia. *Precambrian Res.* 111, 5–30.
- Kellogg, L.H., Hager, B.H., Van der Hilst, R.D., 1999. Compositional stratification in the deep mantle. *Science* 283, 1881–1884.
- Kelly, N.M., Clarke, G.L., Fanning, C.M., 2002. A two-stage evolution of the Neoproterozoic Rayner structural episode; new U–Pb sensitive high resolution ion microprobe constraints from the Oygarden Group, Kemp Land East Antarctica. *Precambrian Res.* 116, 307–330.
- Keppie, J.D., Ortega-Gutierrez, F.O., 1999. Middle American Precambrian basement: a missing piece of the reconstructed 1-Ga orogen. *Geol. Soc. Am. Spec. Pap.* 336, pp. 199–210.
- Keppie, J.D., Ramos, V.A., 1999. Odyssey of terranes in the Iapetus and Rheic oceans during the Paleozoic. *Geol. Soc. Am. Spec. Pap.* 336, pp. 267–276.
- Keppie, J.D., Dostal, J., Cameron, K.L., Solari, L.A., Ortega-Gutierrez, F., Lopez, R., 2003. Geochronology and geochemistry of Grenvillian igneous suites in the northern Oaxacan Complex, southern Mexico: tectonic implications. *Precambrian Res.* 120, 365–389.
- Khudoley, A.K., 1997. Stratigraphic evidence for the Siberia–Laurentia connection and Early Cambrian rifting: comment. *Geology* 25, 570.
- Kirschvink, J.L., 1992. Late Proterozoic low-latitude global glaciation: the snowball earth. In: Schopf, J.W., Klein, C. (Eds.), *The Proterozoic Biosphere*. Cambridge University Press, pp. 51–52.
- Kröner, A., Cordani, U., 2003. African, southern Indian and South American cratons were not part of the Rodinia supercontinent: evidence from field relationships and geochronology. *Tectonophysics* 375, 325–352.
- Kröner, A., Hegner, E., Collins, A.S., Windley, B.F., Brewer, T.S., Razakamanana, T., Pidgeon, R.T., 2000. Age and magmatic history of the Antananarivo Block, central Madagascar, as derived from zircon geochronology and Nd isotopic systematics. *Am. J. Sci.* 300, 251–288.

- Larson, R.L., 1991a. Latest pulse of Earth: evidence for a Mid-Cretaceous super plume. *Geology* 19, 547–550.
- Larson, R.L., 1991b. Geological consequences of superplumes. *Geology* 19 (10), 963–966.
- LeCheminant, A.N., Heaman, L.M., 1994. 779 Ma mafic magmatism in the northwestern Canadian Shield and northern Cordillera: a new regional time-marker. In: *Proceedings of the 8th International Conference. Geochronology, Cosmochronology and Isotope Geology, Program Abstracts*, US Geological Survey Circ. 1107, Berkeley, CA, p. 197.
- Li, Z.X., 2000. New palaeomagnetic results from the “cap dolomite” of the Neoproterozoic Walsh Tillite, northwestern Australia. *Precambrian Res.* 100, 359–370.
- Li, Z.X., 2001. Understanding the precambrian tectonic events in tasmania: clues from south china. *Geol. Soc. Aust. Abst.* 64, 110–111.
- Li, Z.X., Powell, C.M., 2001. An outline of the Palaeogeographic evolution of the Australasian region since the beginning of the Neoproterozoic. *Earth-Sci. Rev.* 53, 237–277.
- Li, Z.X., Zhang, L., Powell, C.M., 1995. South China in Rodinia: part of the missing link between Australia–East Antarctica and Laurentia? *Geology* 23, 407–410.
- Li, Z.X., Zhang, L., Powell, C.M., 1996. Positions of the East Asian cratons in the Neoproterozoic supercontinent Rodinia. *Aust. J. Earth Sci.* 43, 593–604.
- Li, Z.X., Baillie, P.W., Powell, C.M., 1997. Relationship between northwestern Tasmania and east Gondwanaland in the late Cambrian/early Ordovician: Palaeomagnetic evidence. *Tectonics* 16, 161–171.
- Li, Z.X., Li, X.H., Kinny, P.D., Wang, J., 1999. The breakup of Rodinia: did it start with a mantle plume beneath South China? *Earth Planet. Sci. Lett.* 173, 171–181.
- Li, Z.X., Li, X.H., Kinny, P.D., Zhou, H., 2001. Does it take a superplume to break up a supercontinent?—a case for Rodinia. In: *Sircombe, K.N., Li, Z.X. (Eds.), From Basins to Mountains: Rodinia at the Turn of the Century*. *Geol. Soc. Aust., Perth*, pp. 74–77 (Abst. 65).
- Li, X.H., Li, Z.X., Zhou, H., Liu, Y., Kinny, P.D., 2002. U–Pb zircon geochronology, geochemistry and Nd isotopic study of Neoproterozoic bimodal volcanic rocks in the Kangdian Rift of South China; implications for the initial rifting of Rodinia. *Precambrian Res.* 113, 135–154.
- Li, Z.X., Li, X., Zhou, H., Kinny, P.D., 2002. Grenvillian continental collision in South China: new SHRIMP U–Pb zircon results and implications for the configuration of Rodinia. *Geology* 30, 163–166.
- Li, X.H., Li, Z.X., Ge, W.C., Zhou, H.W., Li, W.X., Liu, Y., Wingate, M.T.D., 2003a. Neoproterozoic granitoids in South China: crustal melting above a mantle plume at ca 825 Ma? *Precambrian Res.* 122, 45–83.
- Li, X.H., Li, Z.X., Zhou, H., Liu, Y., Liang, X., Li, W., 2003b. SHRIMP U–Pb zircon age, geochemistry and Nd isotope of the Guandaoshan pluton in SW Sichuan: petrogenesis and tectonic significance. *Sci. China D* 46 (Suppl.), 73–83.
- Li, Z.X., Cho, M., Li, X.H., 2003a. Precambrian tectonics of East Asia and relevance to supercontinent evolution. *Precambrian Res.* 122, 1–6.
- Li, Z.X., Li, X.H., Kinny, P.D., Wang, J., Zhang, S., Zhou, H., 2003b. Geochronology of Neoproterozoic syn-rift magmatism in the Yangtze Craton South China and correlations with other continents: evidence for a mantle superplume that broke up Rodinia. *Precambrian Res.* 122, 85–109.
- Li, Z.X., Evans, D.A.D., Zhang, S., 2004. A 90° spin on Rodinia: possible causal links between the Neoproterozoic supercontinent, superplume, true polar wander and low-latitude glaciation. *Earth Planet. Sci. Lett.* 220, 409–421.
- Li, X.H., Li, W.X., Li, Z.X., Wang, J., 2005. Formation of the South China Block: evidence from Sibaoan orogenic magmatism. In: *Wingate, M.T.D., Pisarevsky, S.A. (Eds.), Supercontinent and Earth Evolution Symposium*. *Geol. Soc. Aus., Fremantle*, p. 86 (Abst. 81).
- Lin, G.C., Li, X.H., Li, W.X., 2007. SHRIMP U–Pb zircon age, geochemistry and Nd–Hf isotope of Neoproterozoic mafic dyke swarms in western Sichuan: Petrogenesis and tectonic significance. *Sci. China D* 50, 1–16.
- Ling, W., Gao, S., Zhang, B., Li, H., Liu, Y., Cheng, J., 2003. Neoproterozoic tectonic evolution of the northwestern Yangtze craton South China: implications for amalgamation and break-up of the Rodinia Supercontinent. *Precambrian Res.* 122, 111–140.
- Loewy, S.L., Connelly, J.N., Dalziel, I.W.D., Gower, C.F., 2003. Eastern Laurentia in Rodinia: constraints from whole-rock Pb and U/Pb geochronology. *Tectonophysics* 375, 169–197.
- Loizenbauer, J., Wallbrecher, E., Fritz, H., Neumayr, P., Khudeir, A.A., Kloetzli, U., 2001. Structural geology, single zircon ages and fluid inclusion studies of the Meatiq metamorphic core complex: implications for Neoproterozoic tectonics in the Eastern Desert of Egypt. *Precambrian Res.* 110, 357–383.
- Lu, S.N., Li, H.K., Zhang, C.L., Niu, G.H., 2008a. Geological and geochronological evidence for Precambrian evolution of the Tarim platform and surroundings. *Precambrian Res.* 160, 94–107.
- Lu, S., Zhao, G., Wang, H., Hao, G., 2008b. Precambrian metamorphic basement and sedimentary cover of the North China Craton: A review. *Precambrian Res.* 160, 77–93.
- Lund, K., Aleinikoff, J.N., Evans, K.V., Fanning, C.M., 2003. SHRIMP U–Pb geochronology of Neoproterozoic Windermere Supergroup, central Idaho; implications for rifting of western Laurentia and synchronicity of Sturtian glacial deposits. *Geol. Soc. Am. Bull.* 115, 349–372.
- Macouin, M., Besse, J., Ader, M., Gilder, S., Yang, Z., Sun, Z., Agrinier, P., 2004. Combined paleomagnetic and isotopic data from the Doushantuo carbonates South China: implications for the “snowball Earth” hypothesis. *Earth Planet. Sci. Lett.* 224, 387–398.
- Malooof, A.C., Halverson, G.P., Kirschvink, J.L., Schrag, D.P., Weiss, B.P., Hoffman, P.F., 2006. Combined paleomagnetic, isotopic and stratigraphic evidence for true polar wander from the Neoproterozoic Akademikerbreen Group, Svalbard. *GSA Bull.* 118, 1099–1124.
- McCabe, C., Van der Voo, R., 1983. Paleomagnetic results from the upper Keweenawan Chequamegon Sandstone: implications for red bed diagenesis and Late Precambrian apparent polar wander of North America. *Can. J. Earth Sci.* 20, 105–112.
- McMenamin, M.A.S., McMenamin, D.L.S., 1990. The Emergence of Animals: The Cambrian Breakthrough, p. 217.
- McWilliams, M.O., McElhinny, M.W., 1980. Late Precambrian paleomagnetism in Australia: the Adelaide Geosyncline. *J. Geol.* 88, 1–26.
- Meert, J.G., 2003. A synopsis of events related to the assembly of eastern Gondwana. *Tectonophysics* 362, 1–40.
- Meert, J.G., Van der Voo, R., 1994. The Neoproterozoic (1000–540 Ma) glacial intervals: no more snowball earth? *Earth Planet. Sci. Lett.* 123, 1–13.
- Meert, J.G., Van der Voo, R., 1996. Paleomagnetic and $^{40}\text{Ar}/^{39}\text{Ar}$ study of the Sinyai Dolerite Kenya: implications for Gondwana assembly. *J. Geol.* 104, 131–142.
- Meert, J.G., Van der Voo, R., Ayub, S., 1995. Paleomagnetic investigation of the Neoproterozoic Gagwe lavas and Mbozi Complex Tanzania and the assembly of Gondwana. *Precambrian Res.* 74, 225–244.
- Mertanen, S., Pesonen, L.J., Huhma, H., 1996. Palaeomagnetism and Sm–Nd ages of the Neoproterozoic diabase dykes in Laanila and Kautokeino, northern Fennoscandia. In: *Brewer, T.S. (Ed.), Precambrian Crustal Evolution in the North Atlantic Region*. *Geol. Soc. London Spec. Pub.* 112, pp. 331–358.
- Metcalfe, I., 1996. Gondwanaland dispersion Asian accretion and evolution of eastern Tethys in Breakup of Rodinia and Gondwanaland and Assembly of Asia. *Aust. J. Earth Sci.* 43, 605–623.
- Metelkin, D.V., Vernikovskiy, V.A., Kazansky, A.Yu., Belonosov, I.V., 2005a. The Siberian Craton in the structure of the Supercontinent Rodinia: analysis of paleomagnetic Data. *Doklady Earth Sci.* 404, 1021–1026.
- Metelkin, D.V., Vernikovskiy, A.V., Kazansky, A.Yu., Bogolepova, O.K., Gubanov, A.P., 2005b. Paleozoic history of the Kara microcontinent and its relation to Siberia and Baltica: Paleomagnetism, paleogeography and tectonics. *Tectonophysics* 398, 225–243.
- Metelkin, D.V., Belonosov, I.V., Gladkochub, D.P., Donskaya, T.V., Mazukabzov, A.M., Stanevich, A.M., 2005c. Paleomagnetic directions from Nersa intrusions of the Biryusa terrane, Siberian Craton, as reflection of tectonic events in the Neoproterozoic. *Russ. Geol. Geophys.* 46, 398–413.
- Mezger, K., Cosca, M.A., 1999. The thermal history of the Eastern Ghats Belt (India) as revealed by U–Pb and $^{40}\text{Ar}/^{39}\text{Ar}$ dating of metamorphic and magmatic minerals; implications for the SWEAT correlation. *Precambrian Res.* 94, 251–271.
- Miller, K.C., Hargraves, R.B., 1994. Paleomagnetism of Some Indian Kimberlites and Lamproites. *Precambrian Res.* 69, 259–267.

- Möller, C., Söderlund, U., 1997. Age constraints on the deformation within the Eastern Segment SW Sweden: Late Sveconorwegian granite dyke intrusion and metamorphic-deformational relations. *Geologiska Föreningen i Stockholm Förhandlingar* 119, 1–12.
- Moore, E.M., 1991. Southwest U.S.–East Antarctic (SWEAT) connection: a hypothesis. *Geology* 19, 425–428.
- Moore, E.M., 2002. Pre-1 Ga (pre-Rodinian) ophiolites: their tectonic and environmental implications. *Geol. Soc. Am. Bull.* 114, 80–95.
- Moore, E.M., Kellogg, L.H., Dilek, Y., 2000. Tethyan ophiolites, mantle convection, and tectonic “historical contingency”: a resolution of the “ophiolite conundrum”. In: Dilek, Y., Moore, E.M., Elthon, D., Nicolas, A. (Eds.), *Ophiolites and Oceanic Crust: New Insights from Field Studies and the Ocean Drilling Program*. Geological Society of America, Boulder, pp. 3–12 (GSA Spec. Pap.).
- Mosher, S., 1998. Tectonic evolution of the southern Laurentian Grenville orogenic belt. *Geol. Soc. Am. Bull.* 110, 1357–1375.
- Murthy, G., Gower, C., Tubrett, M., Pätzold, R., 1992. Paleomagnetism of Eocambrian Long Range dykes and Double Mer Formation from Labrador Canada. *Can. J. Earth Sci.* 29, 1224–1234.
- Natal’in, B.A., 2004. Phanerozoic tectonic evolution of the Chukotka-Arctic block: problems of the rotational model. In: AGU Fall Meet. Suppl., (Abstract GP43C-04).
- Natal’in, B.A., Amato, J.M., Toro, J., Wright, J.E., 1999. Paleozoic rocks of northern Chukotka Peninsula Russian Far East: implications for the tectonics of the Arctic region. *Tectonics* 18, 977–1003.
- Nokleberg, W.J., Parfenov, L.M., Monger, J.W.H., Norton, I.O., Khanchuk, A.I., Stone, D.B., Scotese, C.R., Scholl, D.W., Fujita, K., 2000. Phanerozoic tectonic evolution of the Circum-North Pacific. US Geological Survey Professional Paper 1626, USGS Denver, p. 133.
- Nyman, M.W., Karlstrom, K.E., Kirby, E., Graubard, C.M., 1994. Mesoproterozoic contractional orogeny in western North America: Evidence from ca 1.4 Ga plutons. *Geology* 22, 901–904.
- Occhipinti, S.A., Reddy, S.M., 2005. Mesoproterozoic to Neoproterozoic reworking in the Capricorn orogen Western Australia: evidence from $^{40}\text{Ar}/^{39}\text{Ar}$ dating. *Geol. Soc. Aus. Abst.* 72, 57.
- Onstott, T.C., Hargraves, R.B., 1981. Proterozoic transcurrent tectonics: palaeomagnetic evidence from Venezuela and Africa. *Nature* 289, 131–136.
- Onstott, T.C., Hargraves, R.B., Joubert, P., 1986. Constraints on the evolution of the Namaqua Province II: reconnaissance palaeomagnetic and $^{40}\text{Ar}/^{39}\text{Ar}$ results from the Namaqua Province and the Kheis Belt. *Trans. Geol. Soc. S. Afr.* 89, 143–170.
- Ovchinnikova, G.V., Semikhatov, M.A., Vasil’eva, I.M., Gorokhov, I.M., Kaurava, O.K., Podkovyrov, V.N., Gorokhovskii, B.M., 2001. Pb–Pb age of limestones of the middle Riphean Malgina Formation, the Uchur-Maya region of East Siberia. *Stratigr. Geol. Correl.* 9, 490–502.
- Palmer, H.C., Baragar, W.R.A., Fortier, M., Foster, J.H., 1983. Paleomagnetism of Late Proterozoic rocks, Victoria Island, Northwest Territories, Canada. *Can. J. Earth Sci.* 20, 1456–1469.
- Park, J.K., 1994. Palaeomagnetic constraints on the position of Laurentia from middle Neoproterozoic to Early Cambrian times. *Precambrian Res.* 69, 95–112.
- Park, J.K., Norris, D.K., Larochelle, A., 1989. Paleomagnetism and the origin of the Mackenzie Arc of northwestern Canada. *Can. J. Earth Sci.* 26, 2194–2203.
- Park, J.K., Buchan, K.L., Harlan, S.S., 1995. A proposed giant radiating dyke swarm fragmented by the separation of Laurentia and Australia based on paleomagnetism of ca 780 Ma mafic intrusions in western North America. *Earth Planet. Sci. Lett.* 132, 129–139.
- Patchett, P.J., Bylund, G., 1977. Age of Grenville belt magnetisation: Rb–Sr and palaeomagnetic evidence from Swedish dolerites. *Earth Planet. Sci. Lett.* 35, 92–104.
- Paulsson, O., Andreasson, P.G., 2002. Attempted break-up of Rodinia at 850 Ma: geochronological evidence from the Svea-Kalak Superterrane Scandinavian Caledonides. *J. Geol. Soc.* 159, 751–761.
- Pavlov, V.E., Gallet, Y., Shatsillo, A.V., 2000. Palaeomagnetism of the upper Riphean Lakhanda Group of the Uchur-Maya area and the hypothesis of the late Proterozoic supercontinent. *Fizika Zemli* 8, 23–34 (in Russian).
- Pavlov, V.E., Gallet, Y., Petrov, P.Yu., Zhuravlev, D.Z., Shatsillo, A.V., 2002. Uy series and late Riphean sills of the Uchur–Maya area: isotopic and palaeomagnetic data and the problem of the Rodinia supercontinent. *Geotectonics* 36, 278–292.
- Pease, V., Scott, R., Eliaeson, K., 2005. In: Wingate, M.T.D., Pisarevsky, S.A. (Eds.), *A Baltica Provenance for Detrital Zircons from Early Paleozoic Sediments of Southern Novaya Zemlya Arctic Russia*. *Geol. Soc. Aus. Inc.*, p. 111 (Abs. No. 81).
- Pease, V., Daly, J.S., Elming, S.-Å., Kumpulainen, R., Moczydlowska, M., Puchkov, V., Roberts, D., Saintot, A., Stephenson, R., 2008. Baltica in the Cryogenian, 850–630 Ma. *Precambrian Res.* 160, 46–65.
- Pease, V., Scott, R., Eliaeson, K., 2006. A Baltica province for the Kara terrane. *Bull. Geol. Soc. Finland*, 121.
- Pelechaty, S.M., 1996. Stratigraphic evidence for the Siberia–Laurentia connection and early Cambrian rifting. *Geology* 24, 719–722.
- Pesonen, L.J., Elming, S.A., Mertanen, S., Pisarevsky, S., D’Agrella, M.S., Meert, J.G., Schmidt, P.W., Abrahamsen, N., Bylund, G., 2003. Palaeomagnetic configuration of continents during the Proterozoic. *Tectonophysics* 375, 289–324.
- Pidgeon, R.T., Smith, C.B., Fanning, C.M., 1989. Kimberlite and lamproite emplacement ages in Western Australia. In: Ross, J., et al. (Eds.), *Kimberlites and Related Rocks Volume 1: Their Composition, Occurrence, Origin and Emplacement*. Blackwell Scientific Publications, Carlton, pp. 382–391 (Geol. Soc. Aust. Special Publication).
- Pisarevsky, S.A., Bylund, G., 1998. Palaeomagnetism of a key section of the Proterozoic Zone, southern Sweden. *Geophys. J. Int.* 133, 185–200.
- Pisarevsky, S.A., Bylund, G., 2006. Palaeomagnetism of 935 Ma mafic dykes in southern Sweden and implications for the Sveconorwegian Loop. *Geophys. J. Int.* 166, 1095–1104.
- Pisarevsky, S.A., Natapov, L.M., 2003. Siberia and Rodinia. *Tectonophysics* 375, 221–245.
- Pisarevsky, S.A., Wingate, M.T.D., Harris, L.B., 2003a. Late Mesoproterozoic (ca 1.2 Ga) palaeomagnetism of the Albany–Fraser orogen: no pre-Rodinian Australia–Laurentia connection. *Geophys. J. Int.* 155, F6–F11.
- Pisarevsky, S.A., Wingate, M.T.D., Powell, C.M., Johnson, S., Evans, D.A.D., 2003b. Models of Rodinia assembly and fragmentation. In: Yoshida, M., Windley, B.F., Dasgupta, S. (Eds.), *Proterozoic East Gondwana: Supercontinent Assembly and Breakup*. *Geol. Soc. London Spec. Pub.* 206, pp. 35–55.
- Pisarevsky, S.A., Natapov, L.M., Donskaya, T.V., Gladkochub, D.P., Vernikovsky, V.A., 2008. Proterozoic Siberia: a promontory of Rodinia. *Precambrian Res.* 160, 66–76.
- Pisarevsky, S.A., Wingate, M.T.D., Stevens, M.K., Haines, P.W., 2007. Paleomagnetic results from the Lancer-1 stratigraphic drill hole, Officer Basin, Western Australia, and implications for Rodinia reconstructions. *Aust. J. Earth Sci.* 54, 561–572.
- Powell, C.M., Pisarevsky, S.A., 2002. Late Neoproterozoic assembly of East Gondwana. *Geology* 30, 3–6.
- Powell, C.M., Li, Z.X., McElhinny, M.W., Meert, J.G., Park, J.K., 1993. Paleomagnetic constraints on timing of the Neoproterozoic breakup of Rodinia and the Cambrian formation of Gondwana; with Suppl Data 9335. *Geology* 21, 889–892.
- Powell, C.M., Preiss, W.V., Gatehouse, C.G., Krapez, B., Li, Z.X., 1994. South Australian record of a Rodinian epicontinental basin and its mid-Neoproterozoic breakup (approximately 700 Ma) to form the palaeo-Pacific Ocean. *Tectonophysics* 237, 113–140.
- Powell, C.M., Jones, D.L., Pisarevsky, S., Wingate, M.T.D., 2001. Palaeomagnetic constraints on the position of the Kalahari craton in Rodinia. *Precambrian Res.* 110, 33–46.
- Prave, A.R., 1996. Tale of three cratons: tectonostratigraphic anatomy of the Damara orogen in northwestern Namibia and the assembly of Gondwana. *Geology* 24, 1115–1118.
- Preiss, W.V., 2000. The Adelaide Geosyncline of South Australia and its significance in Neoproterozoic continental reconstruction. *Precambrian Res.* 100, 21–63.
- Radhakrishna, T., Mathew, J., 1996. Late Precambrian (850–800 Ma) palaeomagnetic pole for the south Indian shield from the Harohalli alkaline dykes:

- geotectonic implications for Gondwana reconstructions. *Precambrian Res.* 80, 77–87.
- Rainbird, R.H., de Freitas, T.A., 1997. Stratigraphic evidence for the Siberia–Laurentia connection and Early Cambrian rifting: comment. *Geology* 25, 569–570.
- Rainbird, R.H., Jefferson, C.W., Young, G.M., 1996. The early Neoproterozoic sedimentary Succession B of northwestern Laurentia: correlations and paleogeographic significance. *GSA Bull.* 108, 454–470.
- Rainbird, R.H., Stern, R.A., Khudoley, A.K., Kropachev, A.P., Heaman, L.M., Sukhorukov, V.I., 1998. U–Pb geochronology of Riphean sandstone and gabbro from Southeast Siberia and its bearing on the Laurentia–Siberia connection. *Earth Planet. Sci. Lett.* 164, 409–420.
- Reginussen, H., Ravna, E.J.K., Berglund, K., 1995. Mafic Dykes from Oksfjord, Seiland Igneous Province Northern Norway—geochemistry and paleotectonic significance. *Geol. Mag.* 132, 667–681.
- Renne, P.R., Onstott, T.C., D'Agrella-Filho, M.S., Pacca, I.G., Teixeira, W., 1990. $^{40}\text{Ar}/^{39}\text{Ar}$ dating of 1.0–1.1 Ga magnetizations from the São Francisco and Kalahari cratons: tectonic implications for Pan-African and Brasiliano mobile belts. *Earth Planet. Sci. Lett.* 101, 349–366.
- Rivers, T., 1997. Lithotectonic elements of the Grenville Province: review and tectonic implications. *Precambrian Res.* 86, 117–154.
- Robb, L.J., Armstrong, R.A., Watres, D.J., 1999. The history of granulite-facies metamorphism and crustal growth from single zircon U–Pb geochronology: Namaqualand, South Africa. *J. Petrol.* 40, 1747–1770.
- Ross, G.M., Villeneuve, M.E., 1997. U–Pb geochronology of strange stones in Neoproterozoic diamictites, Canadian Cordillera: implications for provenance and ages of deposition. In: *Radiogenic Age and Isotopic Studies: Report 10. Geol. Surv. Can. Current Res.* 1997-F, pp. 141–155.
- Ross, G.M., Villeneuve, M., 2003. Provenance of the Mesoproterozoic (1.45 Ga) Belt basin (western North America): Another piece of the pre-Rodinian paleogeographic puzzle. *GSA Bull.* 115, 1191–1217.
- Ross, G.M., Parrish, R.R., Winston, D., 1992. Provenance and U–Pb geochronology of the Mesoproterozoic Belt Supergroup (northwestern United States): implications for age of deposition and pre-Panthalassa plate reconstructions. *Earth Planet. Sci. Lett.* 113, 57–76.
- Roy, A.B., 2001. Neoproterozoic crustal evolution of Northwestern Indian Shield: implications on break up and assembly of supercontinents. *Gond. Res.* 4, 289–306.
- Roy, J.L., Robertson, W.A., 1978. Paleomagnetism of the Jacobsville Formation and the apparent polar path for the interval ~1100 to ~670 m.y. for North America. *J. Geophys. Res.* 83, 1289–1304.
- Sadowski, G.R., Bettencourt, J.S., 1996. Mesoproterozoic tectonic correlations between eastern Laurentia and the western border of the Amazon Craton. *Precambrian Res.* 76, 213–227.
- Santosh, M., Iyer, S.S., Vasconcellos, M.B.A., Enzweiler, J., 1989. Late Precambrian alkaline plutons in southwest India: Geochronologic and rare-earth elements constraints on Pan-African magmatism. *Lithos* 24, 65–79.
- Scherstén, A., Årebäck, H., Cornell, D., Hoskin, P., Åberg, A., Armstrong, R., 2000. Dating mafic–ultramafic intrusions by ion-microprobing contact—melt zircon: examples from SW Sweden. *Contrib. Mineral. Petrol.* 139, 115–125.
- Schmidt, P.W., Williams, G.E., 1995. The Neoproterozoic climatic paradox: equatorial palaeolatitude for Marinoan glaciation near sea level in South Australia. *Earth Planet. Sci. Lett.* 134, 107–124.
- Schmidt, P.W., Williams, G.E., Embleton, B.J.J., 1991. Low palaeolatitude of Late Proterozoic glaciation: early timing of remanence in haematite of the Elatina Formation South Australia. *Earth Planet. Sci. Lett.* 105, 355–367.
- Schmidt, P.W., Williams, G.E., Camacho, A., 2005. A revised pole for the ~1070 Ma (?) Alcurra (Kulgera) intrusions. In: Wingate, M.T.D., Pisarevsky, S.A. (Eds.), *Supercontinents and Earth Evolution Symposium 2005*. Geol. Soc. Australia Inc, p. 73 (Abstracts 81).
- Schubert, G., Masters, G., Olson, P., Tackley, P., 2004. Superplumes or plume clusters? *Phys Earth Planet. Inter.* 146, 147–162.
- Sears, J.W., Price, R.A., 1978. The Siberian connection: a case for the Precambrian separation of the North American and Siberian cratons. *Geology* 6, 267–270.
- Sears, J.W., Price, R.A., 2000. New look at the Siberian connection: no SWEAT. *Geology* 28, 423–426.
- Singh, A.K., Vallinayagam, G., 2004. Geochemistry and petrogenesis of anorogenic basic volcanic–plutonic rocks of the Kundal area, Malani Igneous Suite, western Rajasthan India. *Proc. Indian Acad. Sci. (Earth Planet. Sci.)* 113, 667–681.
- Sklyarov, E.V., Gladkochub, D.P., Mazukabzov, A.M., Menshagin, Y.V., Watanabe, T., Pisarevsky, S.A., 2003. Neoproterozoic mafic dike swarms of the Sharyzhalskiy metamorphic massif, southern Siberian craton. *Precambrian Res.* 122, 359–376.
- Smethurst, M.A., Khramov, A.N., Torsvik, T.H., 1998. The Neoproterozoic and Palaeozoic palaeomagnetic data for the Siberian Platform: from Rodinia to Pangea. *Earth Sci. Rev.* 43, 1–24.
- Söderlund, U., Isachsen, C.E., Bylund, G., Heaman, L.M., Patchett, P.J., Vervoort, J.D., Andersson, U.B., 2005. U–Pb baddeleyite ages and Hf, Nd isotope chemistry constraining repeated mafic magmatism in the Fennoscandian Shield from 1.6 to 0.9 Ga. *Contrib. Mineral. Petrol.* 150, 174–194.
- Söderlund, U., Möller, C., Andersson, J., Johansson, L., Whitehouse, M., 2002. Zircon geochronology in polymetamorphic gneisses in the Sveconorwegian orogen SW Sweden: ion microprobe evidence for 1.46–1.42 and 0.98–0.96 Ga reworking. *Precambrian Res.* 113, 193–225.
- Söderlund, U., Patchett, P.J., Vervoort, J.D., Isachsen, C.E., 2004. The ^{176}Lu decay constant determined by Lu–Hf and U–Pb isotope systematics of Precambrian mafic intrusions. *Earth Planet. Sci. Lett.* 219, 311–324.
- Sohl, L.E., Christie-Blick, N., Kent, D.V., 1999. Paleomagnetic polarity reversals in Marinoan (ca 600 Ma) glacial deposits of Australia: Implications for the duration of low-latitude glaciation in Neoproterozoic time. *Geol. Soc. Am. Bull.* 111, 1120–1139.
- Stearn, J.E.F., Piper, J.D.A., 1984. Palaeomagnetism of the Sveconorwegian mobile belt of the Fennoscandian Shield. *Precambrian Res.* 23, 201–246.
- Stein, M., 2003. Tracing the plume material in the Arabian–Nubian Shield. *Precambrian Res.* 123, 223–234.
- Stein, M., Goldstein, S.L., 1996. From plume head to continental lithosphere in the Arabian–Nubian shield. *Nature* 382, 773–778.
- Stern, R.J., 1994. Arc Assembly and Continental Collision in the Neoproterozoic East-African Orogen—implications for the consolidation of Gondwanaland. *Ann. Rev. Earth Planet. Sci.* 22, 319–351.
- Stern, R.J., 2005. Evidence from ophiolites, blueschists, and ultrahigh-pressure metamorphic terranes that the modern episode of subduction tectonics began in Neoproterozoic time. *Geology* 33, 557–560.
- Stump, E., 1995. *The Ross Orogen of the Transantarctic Mountains*. Cambridge University Press, Cambridge, 284 pp.
- Su, Q., Goldberg, S.A., Fullagar, P.D., 1994. Precise U–Pb zircon ages of Neoproterozoic plutons in the Southern Appalachian Blue Ridge and their implications for the initial rifting of Laurentia. *Precambrian Res.* 68, 81–95.
- Sun, S.-S., Sheraton, J.W., 1996. Geochemical and isotopic evolution. In: Glikson, A.Y., et al. (Eds.), *Geology of the Western Musgrave Block, Central Australia, with Particular Reference to the Mafic–Ultramafic Giles Complex*. Aust. Geol. Surv. Org. (AGSO) Bull. 239, pp. 135–143.
- Tackley, P.J., Stevenson, D.J., Glatzmaier, G.A., Schubert, G., 1993. Effects of an endothermic phase transition at 670 km depth in a spherical model of convection in the Earth's mantle. *Nature* 361, 699–704.
- Tanaka, H., Idnurm, M., 1994. Palaeomagnetism of Proterozoic mafic intrusions and host rocks of the Mount Isa inlier Australia: revisited. *Precambrian Res.* 69, 241–258.
- Tassinari, C.C.G., Bettencourt, J.S., Galdes, M.C., Macambira, M.J.B., Lafon, J.M., 2000. In: Cordiani, U.G., Milani, E.J., Thomaz Filho, A., Campos, D.A. (Eds.), *The Amazonian Craton. Tectonic Evolution of South America*, Rio de Janeiro, pp. 41–95.
- Teklay, M., Kröner, A., Mezger, K., 2002. Enrichment from plume interaction in the generation of Neoproterozoic arc rocks in northern Eritrea: implications for crustal accretion in the southern Arabian–Nubian Shield. *Chem. Geol.* 184, 167–184.
- Thorkelson, D.J., 2000. Geology and mineral occurrences of the Slats Creek, Fairchild Lake and “Dolores Creek” areas, Wernecke Mountains, Yukon Territory (106D/16, 106C/13, 106C/14). Exploration and Geological Services Division, Yukon, Indian and Northern Affairs Canada, Bull. 10, 73 pp.
- Thorkelson, D.J., Abbott, J.A., Mortensen, J.K., Creaser, R.A., Villeneuve, M.E., McNicoll, V.J., Lamer, P.W., 2005. Early and Middle Proterozoic evolution of Yukon, Canada. *Can. J. Earth Sci.* 42, 1045–1071.

- Tohver, E., Van der Pluijm, B.A., Van der Voo, R., Rizzotto, G., Scandolaria, J.E., 2002. Paleogeography of the Amazon craton at 1.2 Ga: early Grenvillian collision with the Llano segment of Laurentia. *Earth Planet. Sci. Lett.* 199, 185–200.
- Tohver, E., Van der Pluijm, B.A., Mezger, K., Essene, E.J., Scandolaria, J.E., Rizzotto, G., 2004. Significance of the Nova Brasilândia metasedimentary belt in western Brazil: redefining the Mesoproterozoic boundary of the Amazon craton. *Tectonics* 23, doi: 10.1029/2003TC001563.
- Tohver, E., Van der Pluijm, B.A., Mezger, K., Scandolaria, J.E., Essene, E.J., 2005. Two stage tectonic history of the SW Amazon craton in the late Mesoproterozoic: identifying a cryptic suture zone. *Precambrian Res.* 137, 35–59.
- Torsvik, T.H., Eide, E., 1998. Database of Norwegian geochronology. NGU Report 98-003, 54 pp.
- Torsvik, T.H., Smethurst, M.A., Meert, J.G., VanderVoo, R., McKerrow, W.S., Brasier, M.D., Sturt, B.A., Walderhaug, H.J., 1996. Continental break-up and collision in the Neoproterozoic and Palaeozoic—a tale of Baltica and Laurentia. *Earth-Sci. Rev.* 40, 229–258.
- Torsvik, T.H., Carter, L.M., Ashwal, L.D., Bhushan, S.K., Pandit, M.K., Jamtveit, B., 2001a. Rodinia refined or obscured: palaeomagnetism of the Malani igneous suite (NW India). *Precambrian Res.* 108, 319–333.
- Torsvik, T.H., Ashwal, L.D., Tucker, R.D., Eide, E.A., 2001b. Neoproterozoic geochronology and palaeogeography of the Seychelles microcontinent: the India link. *Precambrian Res.* 110, 47–59.
- Trindade, R.I.F., D'Agrella-Filho, M.S., Epof, I., Brito Neves, B.B., 2006. Paleomagnetism of Early Cambrian Itabaiana mafic dikes (NE Brazil) and the final assembly of Gondwana. *Earth Planet. Sci. Lett.* 244, 361–377.
- Trompette, R., 1994. Geology of Western Gondwana (2000–500 Ma). A.A. Balkema, Rotterdam/Brookfield, p. 350.
- Trompette, R., 1997. Neoproterozoic (~600 Ma) aggregation of Western Gondwana: a tentative scenario. *Precambrian Res.* 82, 101–112.
- Tucker, R.D., Ashwal, L.D., Torsvik, T.H., 2001. U–Pb geochronology of Seychelles granitoids: a Neoproterozoic continental arc fragment. *Earth Planet. Sci. Lett.* 187, 27–38.
- Tyler, I.M., Griffin, T.J., 1990. Structural development of the King Leopold Orogen, Kimberley region, Western Australia. *J. Struct. Geol.* 12, 703–714.
- Tyler, I.M., Pirajno, F., Bagas, L., Myers, J.S., Preston, W.A., 1998. The geology and mineral deposits of the Proterozoic in Western Australia. In: Hodgson, I., Hince, Bernadette (Eds.), *Geology and Mineral Potential of Major Australian Mineral Provinces*. AGSO J. Aust. Geol. Geophys. 17, pp. 223–244.
- Valentine, J.W., Moores, E.M., 1970. Plate-tectonic regulation of animal diversity and sea level: a model. *Nature* 228, 657–659.
- Van der Voo, R., Peinado, J., Scotese, C.R., 1984. Was Laurentia part of an Eocambrian supercontinent? *Geodynamics Ser.* 12, 131–136.
- Van Schmus, W.R., Bickford, M.E., Turek, A., 1996. Proterozoic geology of the east-central Midcontinent basement. In: C.P.A. van der Pluijm B.A. (Ed.), *Basement and Basins of Eastern North America*. GSA Spec. Paper, Geological Society of America, pp. 7–23.
- Veevers, J.J., 2004. Gondwanaland from 650–500 Ma assembly through 320 Ma merger in Pangea to 185–100 Ma breakup: supercontinental tectonics via stratigraphy and radiometric dating. *Earth-Sci. Rev.* 68, 1–132.
- Vernikovsky, V.A., 1997. Neoproterozoic and Late Paleozoic Taimyr orogenic and ophiolitic belts North Asia: a review and models for their formation. In: *Proceedings of the 30th International Geol. Congr. VSP* 7, pp. 121–138.
- Vernikovsky, V.A., Vernikovskaya, A.E., 2001. Central Taimyr accretionary belt (Arctic Asia): Meso-Neoproterozoic tectonic evolution and Rodinia break up. *Precambrian Res.* 110, 127–141.
- Vernikovsky, V.A., Vernikovskaya, A.E., Kotov, A.B., Sal'nikova, E.B., Kovach, V.P., 2003. Neoproterozoic accretion-collisional events on the western margin of the Siberian Craton: new geological and geochronological evidence from the Yenisey Ridge. *Tectonophysics* 375, 147–168.
- Wade, B.P., Barovich, K.M., Hand, M., Scrimgeour, I.R., Close, D.F., 2006. Evidence for Early Mesoproterozoic Arc Magmatism in the Musgrave Block Central Australia: Implications for Proterozoic Crustal Growth and Tectonic Reconstructions of Australia. *J. Geol.* 114, 43–63.
- Walderhaug, H.J., Torsvik, T.H., Eide, E.A., Sundvoll, E.A., Bingen, B., 1999. Geochronology and palaeomagnetism of the Hunnedalen dykes SW Norway: implications for the Sveconorwegian apparent polar wander loop. *Earth Planet. Sci. Lett.* 169, 71–83.
- Wang, J., Li, Z.X., 2003. History of Neoproterozoic rift basins in South China: implications for Rodinia break-up. *Precambrian Res.* 122, 141–158.
- Wang, X.D., Lindh, A., 1996. Temperature-pressure investigation of the southern part of the southwest Swedish Granulite region. *Eur. J. Mineral.* 8, 51–67.
- Wang, X.D., Page, L.M., Lindh, A., 1996. $^{40}\text{Ar}/^{39}\text{Ar}$ geochronological constraints from the southeasternmost part of the eastern segment of the Sveconorwegian orogen: implications for timing of granulite facies metamorphism. *GFF* 118, 1–8.
- Wang, X.L., Zhou, J.C., Qiu, J.S., Zhang, W.L., Liu, X.M., Zhang, G.L., 2006. LA-ICP-MS U–Pb zircon geochronology of the Neoproterozoic igneous rocks from Northern Guangxi South China: Implications for tectonic evolution. *Precambrian Res.* 145, 111–130.
- Warnock, A.C., Kodama, K.P., Zeitler, P.K., 2000. Using thermochronometry and low-temperature demagnetization to accurately date Precambrian paleomagnetic poles. *J. Geophys. Res.* 105, 19435–19453.
- Weber, B., Hecht, L., 2003. Petrology and geochemistry of metaigneous rocks from a Grenvillian basement fragment in the Maya block: the Guichicovi complex, Oaxaca, southern Mexico. *Precambrian Res.* 124, 41–67.
- Weil, A.B., Geissman, J.W., Van der Voo, R., 2004. Paleomagnetism of the Neoproterozoic Chuar Group, Grand Canyon Supergroup, Arizona: implications for Laurentia's Neoproterozoic APWP and Rodinia break-up. *Precambrian Res.* 129, 71–92.
- Weil, A.B., Van der Voo, R., Mac Niocaill, C., Meert, J.G., 1998. The Proterozoic supercontinent Rodinia: paleomagnetically derived reconstructions for 1100 to 800 Ma. *Earth Planet. Sci. Lett.* 154, 13–24.
- Whitehouse, M.J., Windley, B.F., Stoesser, D.B., Al-Khirbash, S., Ba-Bttat, M.A.O., Haider, A., 2001. Precambrian basement character of Yemen and correlations with Saudi Arabia and Somalia. *Precambrian Res.* 105, 357–369.
- Winchester, J.A., 1988. Later Proterozoic environments and tectonic evolution in the northern Atlantic lands. In: Winchester, J.A. (Ed.), *Later Proterozoic Stratigraphy of the Northern Atlantic Regions*. Blackie, Glasgow, pp. 253–270.
- Wingate, M.T.D., 2001. SHRIMP baddeleyite and zircon ages for an Umkondo dolerite sill, Nyanga Mountains, eastern Zimbabwe. *South Afr. J. Geol.* 104, 13–22.
- Wingate, M.T.D., Giddings, J.W., 2000. Age and palaeomagnetism of the Mundine Well dyke swarm Western Australia: implications for an Australia–Laurentia connection at 755 Ma. *Precambrian Res.* 100, 335–357.
- Wingate, M.T.D., Campbell, I.H., Compston, W., Gibson, G.M., 1998. Ion microprobe U–Pb ages for Neoproterozoic basaltic magmatism in south-central Australia and implications for the breakup of Rodinia. *Precambrian Res.* 87, 135–159.
- Wingate, M.T.D., Pisarevsky, S.A., Evans, D.A.D., 2002. Rodinia connections between Australia and Laurentia: no SWEAT, no AUSWUS? *Terra Nova* 14, 121–128.
- Wu, H.C., Zhang, S.H., Li, Z.X., Li, H.Y., Dong, J., 2005. New paleomagnetic results from the Yangzhuang Formation of the Jixian System, North China, and tectonic implications. *Chin. Sci. Bull.* 50, 1483–1489.
- Xiao, S.H., Bao, H.M., Wang, H.F., Kaufman, A.J., Zhou, C.M., Li, G.X., Yuan, X.L., Ling, H.F., 2004. The Neoproterozoic Qurugtagh Group in eastern Chinese Tianshan: evidence for a post-Marinoan glaciation. *Precambrian Res.* 130, 1–26.
- Xu, B., Jian, P., Zheng, H.F., Zou, H.B., Zhang, L.F., Liu, D.Y., 2005. U–Pb zircon geochronology and geochemistry of Neoproterozoic volcanic rocks in the Tarim Block of northwest China: implications for the breakup of Rodinia supercontinent and Neoproterozoic glaciations. *Precambrian Res.* 136, 107–123.
- Yang, Z.Y., Sun, Z.M., Yang, T.S., Pei, Y.L., 2004. A long connection (750–380 Ma) between South China and Australia: paleomagnetic constraints. *Earth Planet. Sci. Lett.* 220, 423–434.
- Young, G.M., 1992. Late Proterozoic stratigraphy and the Canada–Australia connection. *Geology* 20, 215–218.

- Zhang, C.L., Dong, Y.G., Zhao, Y., Wang, A.G., 2003. Geochemistry of Mesoproterozoic volcanics in West Kunlun: evidence for plate tectonic evolution. *Acta Geol. Sin.* 78, 532–542.
- Zhang, S.H., Jiang, G.Q., Zhang, J.M., Song, B., Kennedy, M.J., Christie-Blick, N., 2005. U–Pb sensitive high-resolution ion microprobe ages from the Doushantuo Formation in south China: constraints on late Neoproterozoic glaciations. *Geology* 33, 473–476.
- Zhang, C.L., Li, Z.X., Li, X.H., Ye, H., Wang, A., Guo, K.Y., 2006. Neoproterozoic bimodal intrusive complex in the southwestern Tarim block, northwest China: age, geochemistry and implications for the rifting of Rodinia. *Int. Geol. Rev.* 48, 112–128.
- Zhang, S., Li, Z.X., Wu, H., 2006. New Precambrian palaeomagnetic constraints on the position of the North China Block in Rodinia. *Precambrian Res.* 144, 213–238.
- Zhao, D., 2001. Seismic structure and origin of hotspots and mantle plumes. *Earth Planet. Sci. Lett.* 192, 251–265.
- Zhao, G., Cawood, P.A., Wilde, S.A., Sun, M., 2002. Review of global 2.1–1.8 Ga orogens; implications for a pre-Rodinia supercontinent. *Earth-Sci. Rev.* 59, 125–162.
- Zhao, J.X., Malcolm, M.T., Korsch, R.J., 1994. Characterisation of a plume-related ~ 800 Ma magmatic event and its implications for basin formation in central–southern Australia. *Earth Planet. Sci. Lett.* 121, 349–367.
- Zhou, M.F., Yan, D.P., Kennedy, A.K., Li, Y.Q., Ding, J., 2002. SHRIMP U–Pb zircon geochronological and geochemical evidence for Neoproterozoic arc-magmatism along the western margin of the Yangtze Block South China. *Earth Planet. Sci. Lett.* 196, 51–67.
- Zhou, C.M., Tucker, R., Xiao, S.H., Peng, Z.X., Yuan, X.L., Chen, Z., 2004. New constraints on the ages of Neoproterozoic glaciations in south China. *Geology* 32, 437–440.
- Zonenshain, L.P., Kuzmin, M.I., Natapov, L.M., 1990. *Geology of the USSR: A Plate-Tectonic Synthesis*. Geodynamics Series 21, AGU, Washington, DC, 242 pp.